

PANDEMIC PROVIDER WEEKLY BROWN BAG

Updated April 2023




ARIZONA DEPARTMENT
OF HEALTH SERVICES

Health and Wellness for all Arizonans

Updates


COVID-19 Vaccine Presentations Available to Order in ASIIS Job Aid


- ***NEW*** Novavax has changed from 10-dose multi-dose vials to 5-dose multi-dose vials
 - (NDC 80631-0102-10)
- **Moderna Adult (Ages 12+) - Red Cap** is No Longer Available to Order in ASIIS
 - (NDC 80777-0273-99)
- **Pediatric Moderna (Ages 6-11 years) - Blue Cap with Purple Border on label** is No Longer Available to Order in ASIIS
 - (NDC 80777-0273-99)
- **Janssen** is No Longer Available to Order in ASIIS
- **Moderna Bivalent Booster (Ages 6 months - 5 years) - Dark Pink Cap with Yellow Border on label** has an updated minimum order quantity of 20 doses instead of 100
 - (NDC 80777-0283-99)





COVID-19 Vaccine Presentations Available to Order in ASIIS


Pfizer


Comirnaty (Ages 12+, No Diluent) - Gray Cap

 180-dose minimum order
 • 6-dose multi-dose vial
 • NDC 00069-2025-10


Pfizer-BioNTech Bivalent Booster (Ages 12+, No Diluent) - Gray Cap

 180-dose minimum order
 • 6-dose multi-dose vial
 • NDC 59267-0504-02


Pfizer-BioNTech Bivalent Booster (Ages 12+, No Diluent) - Gray Cap

 50-dose minimum order
 • Single dose vial
 • NDC 59267-1404-02
 • Does not include an ancillary kit

Pediatric Pfizer-BioNTech (Ages 6 months - 5 years) - Blue Cap with Purple Border on label

 100-dose minimum order
 • 10-dose multi-dose vial
 • NDC 59267-1055-04

Pediatric Pfizer-BioNTech (Ages 6 months - 5 years) - Blue Cap with Purple Border on label

 100-dose minimum order
 • 10-dose multi-dose vial
 • NDC 59267-0565-02


Pediatric Pfizer-BioNTech (Ages 6 months - 5 years) - Blue Cap with Purple Border on label

 100-dose minimum order
 • 10-dose multi-dose vial
 • NDC 59267-0078-04


Pediatric Pfizer-BioNTech (Ages 6 months - 5 years) - Blue Cap with Purple Border on label

 100-dose minimum order
 • 10-dose multi-dose vial
 • NDC 59267-0609-02




COVID-19 Vaccine Presentations Available to Order in ASIIS


Moderna

Pediatric Moderna (Ages 6 months - 5 years) - Blue Cap with Magenta Border on label

 100-dose minimum order
 • 10-dose multi-dose vial
 • NDC 80777-0279-99

Moderna Bivalent Booster (Ages 6+) - Blue Cap

 100-dose minimum order
 • 5-dose multi-dose vial
 • NDC 80777-0282-99

Moderna Bivalent Booster (Ages 6 months - 5 years) - Dark Pink Cap with Yellow Border on label

 20-dose minimum order
 • 2-dose multi-dose vial
 • NDC 80777-0283-99

Novavax

Novavax (Ages 12+)

 50-dose minimum order
 • 5-dose multi-dose vial
 • NDC 80631-0102-10

602-364-3899
 ASIISHelpDesk@azdhs.gov
<https://azdhs.gov/vfc>

ADHS Vaccine Reference Sheets - Pfizer-BioNTech

Pediatric Pfizer **Monovalent & Bivalent** (Ages 6mo - 4 years) - **Maroon Cap**

- [PEDIATRIC Pfizer Vaccine Reference Sheet](#)

Pfizer **Bivalent Booster** (Ages 5-11) - **Orange Cap**

- [Pfizer Vaccine Reference Sheet](#)

Pfizer (Ages 5-11) - **Orange Cap**

- [PEDIATRIC Pfizer Vaccine Reference Sheet](#)

Comirnaty/Pfizer (Ages 12+) - **Gray Cap**

- [Comirnaty/Pfizer Vaccine Reference Sheet](#)

Pfizer **Bivalent Booster** (Ages 12+, No Diluent) - **Gray Cap**

- [Pfizer Vaccine Reference Sheet](#)

Pfizer-BioNTech Bivalent Booster
(Ages 12+, No Diluent) - Gray Cap

Vaccine Storage Pfizer-BioNTech's COVID-19 Vaccine (Gray Cap) Ages 12+

Vaccine Storage Pfizer-BioNTech's Pediatric COVID-19 Vaccine (Orange Cap) Ages 5-11

Pfizer-BioNTech Bivalent Booster
(Ages 5 - 11) - Orange Cap

Pfizer-BioNTech's Pediatric COVID-19 Vaccine (Maroon Cap) Monovalent & Bivalent
Ages 6 Months through 4 years

Ultra-Cold Temperature Freezer

- -90.0°C to -60.0°C
- Can be stored for 18 months at ultra-low temps from manufactured date

Thermal Shipping Container (DO NOT USE FOR STORAGE)

- Pfizer-BioNTech's Pediatric COVID-19 Vaccine (Maroon Cap) CANNOT be stored in the thermal shipping container

Freezer (DO NOT USE FOR STORAGE)

- The Pfizer BioNTech Pediatric COVID-19 Vaccine (Maroon Cap) CANNOT be stored in the freezer

Refrigerator

- 2.0°C to 8.0°C
- Store undiluted vaccine vials for up to 10 weeks

ADHS Vaccine Reference Sheets - Moderna

Moderna **Bivalent Booster** (Ages 6 months - 5 years) - **Dark Pink Cap** with **Yellow Border** on label

- [Moderna Vaccine Reference Sheet](#)

Pediatric Moderna **Blue Cap** with **Magenta border** on label (Ages 6mo - 5 years)

- [Moderna Vaccine Reference Sheet](#)

Pediatric Moderna **Blue Cap** with **Purple Border** on label (Ages 6 - 11 years)


- [Moderna Vaccine Reference Sheet](#)

Moderna **Bivalent Booster (Ages 6+) - Blue Cap**

- [Moderna Vaccine Reference Sheet](#)

Moderna (Ages 12+)

- [Moderna Vaccine Reference Sheet](#)




COVID-19 Vaccination Schedule for Primary and Booster Doses

COVID-19 Vaccination Schedule for people who are **NOT** and who **ARE** Moderately or Severely Immunocompromised



COVID-19 Vaccination Schedule for people who are **NOT** Moderately or Severely Immunocompromised


People ages 6 months through 4 years




People age 5 years



People ages 6 through 11 years



People ages 12 years and older



People ages 18 years and older who previously received Janssen primary series dose*



*For people who previously received a monovalent booster dose(s), the bivalent booster dose is administered at least 2 months after the last monovalent booster dose.

† A monovalent Novavax booster dose maybe used in limited situations in people 18 years and older who completed a primary series using any COVID-19 vaccine, have not received any previous booster dose(s), and are unable or unwilling to receive an mRNA vaccine. The monovalent Novavax booster dose is administered at least 6 months after completion of a primary series.

‡ Janssen COVID-19 dose should only be used in certain limited situations. See: <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix-a.html#appendix-a>




COVID-19 Vaccination Schedule for people who **ARE** Moderately or Severely Immunocompromised

People ages 6 months through 4 years



People age 5 years



People ages 6 through 11 years



People ages 12 years and older




People ages 18 years and older who previously received Janssen primary series dose*



Monoclonal antibodies (EVUSHELD™) for COVID-19 pre-exposure prophylaxis

People ages 12 years and older (must weigh at least 40kg)



*For people who previously received a monovalent booster dose(s), the bivalent booster dose is administered at least 2 months after the last monovalent booster dose.

† A monovalent Novavax booster dose maybe used in limited situations in people 18 years and older who completed a primary series using any COVID-19 vaccine, have not received any previous booster dose(s), and are unable or unwilling to receive an mRNA vaccine. The monovalent Novavax booster dose is administered at least 6 months after completion of a primary series.

‡ Janssen COVID-19 dose should only be used in certain limited situations. See: <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix-a.html#appendix-a>

***For people who previously received a monovalent booster dose(s), the bivalent booster dose is administered at least 2 months after the last monovalent booster dose.**

† A monovalent Novavax booster dose maybe used in limited situations in people 18 years and older who completed a primary series using any COVID-19 vaccine, have not received any previous booster dose(s), and are unable or unwilling to receive an mRNA vaccine. The monovalent Novavax booster dose is administered at least 6 months after completion of a primary series.

‡ Janssen COVID-19 dose should only be used in certain limited situations. See:


<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix-a.html#appendix-a>


COVID-19 Vaccine Program

ADHS COVID-19 Vaccine Website


azhealth.gov/covid19vaccine

COVID-19 Vaccines




 Coronavirus Disease 2019 (COVID-19) Announcements


[Back to Everyone](#)




General Vaccine Info




Provider Resources




Find COVID-19 Vaccines




Vaccine Registration




Patient Portal



VAPAC




Vaccine FAQs



County Vaccine Resources

COVID-19 Menu

COVID-19 

Home

Vacunas en Español

General Vaccine Info

Provider Resources

Find COVID-19 Vaccines

Vaccine Registration

Patient Portal


VAPAC

Vaccine FAQs

County Vaccine Resources

ADHS Find Vaccine webpage azhealth.gov/findvaccine

- The locations of vaccination sites
- Filter to sort by vaccine type
- Use the patient portal at podvaccine.azdhs.gov to make an appointment for a relative
- Appointments recommended but no longer required at state-operated sites



Pandemic Provider Onboarding Start Form

- All facilities must onboard to be able to order/administer COVID-19 vaccines
- [Pandemic Vaccine Provider Onboarding Tool](#) - Job Aid (bottom right)
- Wait until you're at the facility to onboard so you can upload photos of inside and outside of units
- Read through forms for important requirements
- Note: Per the CDC Agreement, COVID-19 vaccines must be recorded in the vaccine recipient's record, the required information reported to the relevant state, local, or territorial public health authority, and decremented from the inventory **within 24 hours of administration.**

ARIZONA DEPARTMENT OF HEALTH SERVICES

Pandemic Vaccine Provider Onboard Form 1

The Arizona Immunization Program Office (AIPO) does not know if/when COVID-19 vaccines will become available. However we want to be prepared and engage our partners now, so we are ready when vaccines are available for distribution in Arizona.

Both VFC and non-VFC providers who would like to administer future COVID-19 vaccines must complete the Pandemic Provider Onboarding survey forms.

After submitting this form, you will be sent to a page with a series of subsequent survey forms. The forms serve two purposes: 1) to ensure the signatory provider knows the requirements, and 2) to ensure the facility is able to meet each requirement. The signatory provider is required to complete all of the surveys and be approved by the AIPO before this facility will be able to order future potential pandemic vaccines.

As we learn more about future vaccines we may add additional survey forms to the onboarding tool to pass the information along.

Onboarding is not an instantaneous process. It will take AIPO time to add providers to ASIS and review requirements. Onboard now. Plan for it to take time.

The onboarding tool has the high level requirements for the signatory provider. Provider staff can go to AIPO Train to learn how to order, receive, store, administer, document, and account for pandemic vaccines in ASIS.

We appreciate your continued efforts to help Arizonans be healthy and vaccinated.

Click the link to view the onboard job aid to help you complete the survey forms. You are currently on Form 1.
Please contact our help desk if you have further questions: (602) 364-3829

ARIZONA DEPARTMENT OF HEALTH SERVICES

Pandemic Vaccine Provider Onboarding Tool

How to get started

1. Click the onboarding link [Pandemic Vaccine Provider Onboard Form 1](#) and fill out the form
2. Click submit - this will take you to survey queue page
3. Continue to complete other surveys that are not marked "Complete". A link to this page will be emailed to you

Contact and Shipping Information

- Facility location information
- Signatory provider info (title, license, NPI)
- Primary and backup vaccine coordinator information
- Not sure if you're a VFC/VCA provider? Select no when it asks if you are a VFC provider; this is not necessary to participate

Storage and Handling

- Photos of your cold storage units showing the inside of the units
- Brand and model of each cold storage unit
- Data logger usage info
- Read requirements

Arizona State Immunization Information System (ASIS)

- Are you currently entering/transferring immunization data into ASIS?
- Read requirements

Vaccine Planning

- Read through content so you can plan and be prepared

CDC Agreement Section A

- CMO & CEO signatories
- For organizations: follow the instructions for Section A in FAQs. Follow the decision tree on the final page to determine whether you need to follow organization instructions

CDC Agreement Section B

- Facility type
- Populations served
- Storage unit capacity
- Must be signed by the signatory provider/the primary vaccine coordinator

Prescribing Providers (part of CDC Agreement)

- Submit multiple times - one for each prescribing provider
- Enter each prescriber's name, title, and license number

Revised November 2020

FAQs

I clicked "Save and Return" and lost my place. What do I do?

- Find the email with the link to the survey queue page
- Click the survey queue page link
- Click on the survey you were working on
- Enter return code
- Continue survey
- If you did not save a return code, click the "start over" button to start the survey form again. You will not lose previously submitted survey forms

How will I know if I'm done?

- On the survey queue page, surveys marked "Complete" are done and surveys with a "Begin Survey" button are incomplete.

What training do I need?

- The [AIPO TRAIN](#) has training courses for staff who are responsible for daily tasks, including:
 - » How to order, receive, and account for doses in ASIS
 - » How to use data loggers
 - » Storage & handling requirements
- Facilities are responsible for knowing and following the guidelines in the AIPO TRAIN training modules

What are the instructions for group organizations with several facility locations?

- The CDC Provider Agreement has Section A and Section B
- CMO & CEO for organizations must complete and sign Section A only once for all facility locations within the organization
- Select one facility location whose survey form will be used
- Inform the other facilities which facility location was selected
- Share the link for the survey with the CMO & CEO
- Use the "Sign and return later" button to allow both the CMO & CEO to sign the same survey form
- After the Section A survey form is signed and submitted by the CMO & CEO, a code number will be emailed to the CMO & CEO
- Have the CMO & CEO forward the email with the code number to all of the organization's facility locations
- The other facilities will enter the code number that was forwarded to them into the survey form titled "Enter code" from section A
- Complete all other sections of the CDC Agreement and onboard survey forms

NOTE: Facilities that are part of an Organization must ensure that their CMO & CEO complete a section A survey form for their Organization. No COVID-19 vaccines will be shipped to facilities whose organization has not completed/signed section A

COVID vaccinators will need to be enrolled with AHCCCS in order to be able to bill the vaccine administration fee for AHCCCS beneficiaries. Providers can register with AHCCCS at: <https://www.azahcccs.gov/PlansProviders/APPE/Access.html>

Revised November 2020

Required Reporting

- ASIIS
 - for inventory accounting and dose administration data within 24 hours
 - Required even if you use an EHR, ADHS VMS app, etc.
 - Verify ASIIS Lot Number > Reconciliation page (ASIIS Inventory) numbers should be matched between the quantity on hand and physical inventory columns
 - Vaccine Inventory Management course in AIPO Train
- Vaccines.gov/CDC VaccineFinder weekly inventory by Fridays
- County surveys as required by your local jurisdiction

Your ASIIS inventory is used by local, state, and federal leaders to make vaccine ordering decisions - It MUST be accurate

COVID-19 Inventory Management Reminders

In order to ensure that provider sites are being good stewards of the COVID-19 vaccines, the CDC has emphasized that each COVID-19 vaccination site should have at most 3-4 weeks of vaccine inventory on hand.

- Place smaller, more frequent orders to avoid stockpiling doses
- Use the data in ASIIS to determine how many doses you use & how many doses are on hand to place an order for an appropriate amount of doses
- Use the [ADHS vaccine transfer matchmaker website](#) to request a smaller quantity of doses then can be ordered in ASIIS
- Administered doses must be entered in ASIIS within 24 hours and decremented from the ASIIS inventory
- [CDC VaccineFinder \(vaccines.gov\)](#) inventory must be updated weekly by Friday
- Wasted/Expired Doses must be reported on signed and complete [wasted/expired forms](#) weekly

Off-Label Use

Pandemic providers should be aware that they must use COVID-19 vaccines according to FDA and CDC guidance. Use of COVID-19 vaccines outside of FDA and CDC recommendations (“off label”) is a violation of the provider agreement.

- Providers may not be covered under PREP Act and therefore not have the immunity from prosecution that the PREP Act provides.
- Recipients of an “off-label” COVID-19 vaccine dose may not be covered by the Countermeasure Injury Compensation Program if they were to have serious adverse events from the vaccine.
- Providers who violate the CDC agreement may not be able to remain as part of CDC programs.
- COVID-19 vaccine administration fees may not be covered if the dose were given “off-label.”

CDC Website: VFC vs. COVID-19 Vaccination Programs

- Separate programs with separate agreements
- Distinct requirements for each
- Help VFC providers understand difference between program requirements

Vaccines & Immunizations

CDC > COVID-19 Vaccination > Provider Requirements and Support

- COVID-19 Vaccination
 - Product Info by U.S. Vaccine
 - Clinical Care
 - Provider Requirements and Support**
 - How to Enroll as a Healthcare Provider
 - Inventory Management Best Practices
 - Vaccines for Children Program vs. CDC COVID-19 Vaccination Program**
 - Training and Education
 - Vaccine Recipient Education
 - Health Departments
 - Planning & Partnerships
 - Vaccine Effectiveness Research
 - Vaccination Toolkits

Vaccines for Children Program vs. CDC COVID-19 Vaccination Program

As Emergency Use Authorization of COVID-19 vaccine products expand to include adolescents and children, it is critical to enroll providers in the COVID-19 Vaccination Program to ensure equitable access to COVID-19 vaccination services. Providers enrolled in the Vaccines for Children (VFC) program are well situated to serve in this capacity due to their direct access to the younger patient population and their familiarity with vaccine administration and federal vaccine programs. Though the VFC and COVID-19 Vaccination programs are both federal government programs, they each have distinct requirements based on the associated funding legislation. For this reason, the provider agreements remain separate, and VFC providers must sign and adhere to the requirements of the *CDC COVID-19 Vaccination Program Provider Agreement* in order to receive and administer COVID-19 vaccines. The table below has been developed to assist VFC providers in understanding the difference in the programs' requirements. **Program differences are in bold.**

	VFC Program	COVID-19 Vaccination Program
Provider Enrollment	<ul style="list-style-type: none">Providers enroll via state/local immunization program enrollment system and procedures.Providers must complete and sign state/local immunization program Vaccines for Children Program Provider Agreement and VFC Program Provider Profile Form.	<ul style="list-style-type: none">Providers enroll via state/local immunization program enrollment system and procedures.Providers must complete and sign CDC COVID-19 Vaccination Program Provider Agreement, Sections A and B.
Vaccine Ordering	<ul style="list-style-type: none">Providers order routine childhood vaccines via state/local immunization program-designated ordering system and procedures.	<ul style="list-style-type: none">Providers order COVID-19 vaccines via state/local immunization program-designated ordering system and procedures.Providers must be fully trained in vaccine


Vaccines.gov/CDC VaccineFinder Provider Resources

The “organization email” listed in Section A of the CDC Agreement will get an email from vaccinefinder@auth.castlighthealth.com to sign up

- Weekly reporting by Friday of on-hand inventory quantities is a requirement
- Activating a location to display to the public is optional
- Providers can update display to the public function at any time
- Once COVID-19 vaccine locations are launched in Vaccines.gov, changes show publicly within 24 hours
- Check phone number in provider portal before making site public
- [Vaccines.gov](https://www.vaccines.gov) homepage - patients can find vaccines
- NOTE: If a site that is set to display to the public fails to update inventory within 72 hours, the availability information for each vaccine will display as “Call to confirm” on Vaccines.gov.

COVID-19 Administration Reporting Systems (CARS) Help Desk

- Monday through Friday, 8:00 am to 8:00 pm ET.
- CARS_HelpDesk@cdc.gov
- 1-833-748-1979
- Provide PIN when calling




Changes to the Required Temperature Incident Reporting Process for COVID-19 Vaccine Temperature Incidents

The Bureau of Immunization Services (BIZS) is introducing changes to the required temperature incident reporting process for COVID-19 vaccine temperature incidents. This process is intended to improve incident turnaround times for providers by allowing them to take ownership of the process by following the steps in the [Temperature Incident Instructions Checklist](#) to obtain a viability determination from the manufacturer.

This is a required process for reporting all temperature incidents involving COVID-19 vaccines. The BIZS also encourages you to please fill out the survey at the end of the process as we value your feedback.

Incident Report Checklist



<https://redcapaipo.azdhs.gov/surveys/?s=AY3C74XEPTWDXI4H>
ArizonaVEC@azdhs.gov
602.364.3642

Temperature Incident Instructions

- Isolate the vaccines in the affected unit, label them DO NOT USE and keep them in the affected unit. Do not administer any doses exposed to out of range temperatures.
- Download the data logger report(s) from the data logger monitoring the affected unit(s).
- Determine how long the unit was out of appropriate temperature range and the highest and/or lowest out of range temperature documented on the data logger report.
 - If any vaccines in the affected unit have been involved in a previous temperature incident, you will need to include the time of the previous incident and the highest/lowest out of range temperature the vaccines were exposed to in your report to the manufacturer(s).

Appropriate temperature ranges:

- Refrigerator: 2.0°C to 8.0°C OR 36.0°F to 46.0°F
- Standard Freezer: -50.0°C to -15.0°C OR -58.0°F to 5.0°
- Ultra Low Temp Freezer: -130.0°F to -76.0°F OR -90.0°C to -60.0°C

- Create a list of all affected vaccines - include the type of vaccine, the lot numbers, the expiration dates and the manufacturers.
- Contact each manufacturer - provide them with the information you obtained from the data logger reports (length of time vaccines were exposed to out of range temperatures and the highest and/or lowest temperature recorded), as well as affected vaccine information, and any previous incident information as requested.
- If reporting the incident to the manufacturer via phone or through email - request a statement of viability be emailed to you by the manufacturer(s). If you are able to check the viability online with an excursion tool from the manufacturer, download a copy of the viability statement for ALL affected vaccines.
- Report the outcome of the incident to the AIPO at this link: <https://redcapaipo.azdhs.gov/surveys/?s=AY3C74XEPTWDXI4H>
- If the affected vaccines are deemed viable and the information reported to the manufacturers and the AIPO was accurate, you may resume use of viable vaccines once the incident has been reported to the AIPO using the above link in step 7.

ts readily available to

time of the previously
d unit has gone back

ident. If the data logger
e more data logger
riate temperature range.

data logger report -
in.

piration date for all).

Information

er Pediatric
urple Cap
Age 12+)

) 438-1985

Daily COVID-19 Vaccine Tasks

Temperature Monitoring

- Twice per day monitor temperatures using the [approved data logger](#)
 - Document that you monitored temperatures using the [paper temp log](#)
 - Comirnaty/Pfizer ULT [Celsius](#) [Fahrenheit](#)
 - Refrigerator [Celsius](#) [Fahrenheit](#)
 - Moderna Freezer [Celsius](#) [Fahrenheit](#)
 - Document current, min, max, time, and initials
- **If there are out of range temperatures, stop using the vaccines and submit an [incident report](#) to the BIZS**
- Twice per month download and save the data logger data reports
 - Keep the data logger reports readily available for 6 yearsSubmit the data logger reports (in an acceptable file format: .xls, .txt, .ltd or .csv) to the BIZS upon request

Take a physical inventory count of doses in the cold storage units

- Compare the physical count to ASIIS lot number reconciliation inventory
- The inventories should be the exact same if doses given are entered properly in ASIIS. If you need to troubleshoot, use this [job aid](#).
- Enter the weekly inventory by Fridays into CDC Vaccines.gov

Document doses administered in ASIIS within 24 hours

- Shipments must be marked “received” in ASIIS prior to administration in order for the doses to decrement from the inventory

If you would no longer like to participate as a pandemic provider please follow these steps

1. Account for all doses in ASIIS

- a. You are responsible for accounting for ALL of the doses shipped to you
- b. Ensure all doses shipped to the facility were received in ASIIS
- c. Ensure all doses administered were decremented from the ASIIS inventory
- d. ASIIS inventory reconciliation screen should be accurate

2. If you still have doses

- a. The doses will need to be transferred to another COVID-19 provider
- b. Transfers must have prior approval in ASIIS

3. Enter zero in CDC Vaccines.gov inventory when your inventory is depleted

4. Notify ADHS *after* doses have been accounted for

AIPO Train

- Training on ordering, receiving, and accounting for doses in ASIIS
- Information on data loggers - setting up, downloading data
- Onboarding resources
- Mass immunizations

How to Register

1. Go to <https://AIPO.myabsorb.com/?KeyName=PandemicProviders> (This link automatically enters the enrollment key)
2. Enter required fields
3. Select Sign up
4. Click email activation link (important)

How to Log In

1. Go to <https://AIPO.myabsorb.com/#/login>
2. Enter username and password (username will be email address)
3. Select Log In

For existing AIPO Train users: Enter enrollment key “PandemicProviders” to access pandemic-specific training

Questions? Email AIPOTrain@azdhs.gov

Administration Fee

- CMS has a toolkit to help health care providers prepare to administer vaccines
- Vaccine doses given to the American people at no cost
- Providers contractually agree to administer a COVID-19 vaccine regardless of an individual's ability to pay and regardless of their coverage status
- May not seek any reimbursement, including through balance billing, from a vaccine recipient
- People without health insurance can get COVID-19 vaccine at no cost
- If you give COVID-19 vaccines to patients who are covered by **AHCCCS** you will need to be enrolled with AHCCCS in order to bill AHCCCS the administration fee
 - [Register with AHCCCS here](#)

Facilities cannot bill patients for the cost or admin fee for COVID-19 vaccines.

CVX Code and CPT Code Resources

- CMS - [Helpful for Billing](#)
- CDC - [Helpful for Uploading Data in EHR and ASIIS](#)
- ADHS - [Arizona HL7 Specific Rules](#)

TAPI - The Arizona Partnership for Immunizations (State immunization coalition)

- [COVID-19 Resources](#)
- [COVID-19 Vaccine Billing Policy Information](#)
- [Resources for immunizations at off-site locations](#)

CDC Guidance for Planning Vaccination Clinics Held at Satellite, Temporary, or Off-Site Locations

CDC has issued revised [Guidance for Planning Vaccination Clinics Held at Satellite, Temporary, or Off-Site Locations](#) to assist with jurisdictional planning and implementation of satellite, temporary, or off-site vaccination clinics by public and private vaccination organizations.

The guidance is broken down into four categories:

- [Planning activities](#)
- [Pre-clinic activities](#)
- [During the clinic activities](#)
- [Post-clinic activities](#)

The guidance also provides information on additional considerations required during the COVID-19 pandemic, including physical distancing, personal protective equipment (PPE), and enhanced sanitation efforts

Training for Healthcare Professionals

- [CDC COVID-19 Vaccine Training Modules](#)
- [Public Health Foundation Training Plan](#)

 Centers for Disease Control and Prevention
CDC 24/7: Saving Lives, Protecting People™

Search 

[A-Z Index](#)

[Advanced Search](#)

Immunization Education & Training





COVID-19 Vaccine Training Modules


 [COVID-19 Vaccine Training: General Overview of Immunization Best Practices for Healthcare Providers](#)

 [Janssen COVID-19 Vaccine \(Johnson & Johnson\): What Healthcare Professionals Need to Know](#) (Updated May 18, 2021)


 [Moderna COVID-19 Vaccine: What Healthcare Professionals Need to Know](#) (Updated May 18, 2021)





 [Pfizer-BioNTech COVID-19 Vaccine: What Healthcare Professionals Need to Know](#) (Updated May 24, 2021)

Page last reviewed: May 24, 2021



HOME COURSE CATALOG CALENDAR RESOURCES HELP

Search 

Name	Completed Date	Score	Hours	Status
Vaccine Storage and Handling 				
Immunization: You Call the Shots-Module Ten-Storage and...			1h	
Keys to Storing and Handling Your Vaccine Supply-2018 (...)			0.9h	
Epidemiology and Prevention of Vaccine-Preventable Dise...			1h	
Vaccine Administration 				
E-LEARN: VACCINE ADMINISTRATION (You Call the Shot...			1h	
Vaccine Administration Videos			1h	
Communicating with Patients about Vaccines 				
How Nurses and Medical Assistants Can Foster a Culture...			0.75h	
"#HowIRecommend" Vaccination Video Series			0.5h	
Epidemiology and Prevention of Vaccine-Preventable Dise...			1h	
COVID-19 Vaccine Training and Clinical Materials 				
COVID-19 Vaccine Training: General Overview of Immuniz...			0.5h	
Pfizer-BioNTech COVID-19 Vaccine: What Healthcare Prof...			0.5h	
Moderna COVID-19 Vaccine: What Healthcare Professiona...			0.5h	
Janssen COVID-19 Vaccine (Johnson & Johnson): What H...			0.5h	

Immunization Action Coalition - immunize.org

- Info on administering vaccines
- Handouts for staff and patients
- “Ask the Experts” page
 - Includes [COVID-19 vaccines](#)
- Option to sign up for IAC Express email
 - Weekly email with updates to information and resources

The screenshot shows the homepage of the Immunization Action Coalition (IAC) website. The header includes navigation links: HOME, ABOUT IAC, CONTACT, A-Z INDEX, DONATE, SHOP, and SUBSCRIBE. There are also links for 'For Healthcare Professionals', 'For the Public', and 'For Coalitions'. The main title 'Immunization Action Coalition' is prominently displayed, followed by a 'Sign up for email newsletter' button and a search bar. Below the header, a horizontal menu lists categories: Favorites, Handouts & Staff Materials, Clinic Tools, Vaccine Information Statements, Vaccines, and Talking about Vaccines. The main content area features a 'Welcome' sidebar with links to IAC Express, Ask the Experts, Handouts, Immunization News, and Shop IAC. The central section highlights 'Handouts' with a call to action 'Free educational materials for patients and staff' and a box stating 'Over 250 materials from IAC experts ready for your use', with links to 'Handouts Main Page Categorized' and 'View All Materials Sorted alphabetically'. Below this, there's a section for 'IAC Express' describing it as a weekly newsletter and a 'SUBSCRIBE' button. To the right, a 'WEBINAR WITH CDC SPEAKERS' section promotes a 'WATCH' video titled 'Updates on CDC's Safety Monitoring and Communications for COVID-19 Vaccines'. Further down, a 'Vaccinating Adults: A Step-by-Step Guide' is available for download as a 'FREE 142-page Guide'. The bottom section includes 'IAC Publications' with links to 'IAC Express - Email news', 'Needle Tips Archive', and 'Vaccinate Adults Archive', and an 'Ask the Experts!' section with a call to action 'Read Ask the Experts'. On the far right, there are several promotional banners, including one for 'IAC's COVID-19 web page', another for 'How to Vaccinate Children, Teens, and Adults During the Time of Covid-19', and a 'WATCH' section for a documentary on IAC's 30-year history titled 'Protecting Health: Saving Lives'. Social media follow links for Facebook, Twitter, and YouTube are also present, along with buttons to 'Read IAC Express', 'Subscribe - It's Free', and 'Make a Donation'.

CDC COVID-19 Vaccination Clinical Resources

COVID-19 Vaccination

Clinical Resources for Each COVID-19 Vaccine

Find information for COVID-19 vaccination administration, storage and handling, reporting, and patient education for each specific vaccine

[Product Information by US Vaccine](#)



ACIP
Recommendations



Storage and
Handling



General Vaccine
Administration



Training and
Education



V-safe



Clinical Considerations



Emergency Use
Authorizations (EUAs)



Vaccination Provider
Requirements &
Support



Vaccination Data &
Reporting Systems




Planning &
Partnerships

Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Authorized in the United States



Other resources:

- [Summary Document for Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Authorized in the United States](#)
- [COVID-19 Vaccine Administration Errors and Deviations](#)



Summary Document for Interim Clinical Considerations

for Use of COVID-19 Vaccines Currently Authorized or Approved in the United States

COVID-19 vaccine products currently approved or authorized in the United States

Pfizer-BioNTech								
Age indication	Vaccine composition	Vaccine vial cap color	Label border color	Dilution required	Primary series		Booster doses	
					Dose	Injection volume	Dose	Injection volume
6 months–4 years	Monovalent (Use for 1st and 2nd Dose)	Maroon	Maroon	Yes	Doses 1 and 2: 3 µg/0.2 mL		A booster dose is not authorized for children who received a 3-dose primary series regardless which vaccine (monovalent or bivalent) was administered for Dose 3.	
6 months–4 years	Bivalent (Use for 3rd Dose)	Maroon	Maroon	Yes	Dose 3: 3 µg/0.2 mL			
5–11 years	Monovalent	Orange	Orange	Yes	10 µg	0.2 mL	NA	NA
5–11 years	Bivalent	Orange	Orange	Yes	NA	NA	10 µg	0.2 mL
12 years and older	Monovalent	Gray	Gray	No	30 µg	0.3 mL	NA	NA
12 years and older	Bivalent	Gray	Gray	No	NA	NA	30 µg	0.3 mL
Moderna								
Age indication	Vaccine composition	Vaccine vial cap color	Label border color	Dilution required	Primary series		Booster doses	
					Dose	Injection volume	Dose	Injection volume
6 months–5 years	Monovalent	Dark blue	Magenta	No	25 µg	0.25 mL	NA	NA
6 months–5 years	Bivalent*	Dark pink	Yellow	No	NA	NA	10 µg	0.2 mL
6–11 years	Monovalent	Dark blue	Purple	No	50 µg	0.5 mL	NA	NA
6–11 years	Bivalent	Dark blue	Gray	No	NA	NA	25 µg	0.25 mL
12 years and older	Monovalent	Red	Light blue	No	100 µg	0.5 mL	NA	NA
12 years and older	Bivalent	Dark blue	Gray	No	NA	NA	50 µg	0.5 mL

CDC Vaccine Storage and Handling Toolkit

✉ Get Email Updates

To receive email updates about this page, enter your email address:

[What's this?](#)

Submit

Related Links

[Vaccines & Immunizations](#)

View or Print Toolkit

The Vaccine Storage and Handling Toolkit is a comprehensive guide that reflects best practices for vaccine storage and handling from Advisory Committee on Immunization Practices (ACIP) recommendations, product information from vaccine manufacturers, and scientific studies.

Vaccine Storage and Handling Resources

Access additional [resources](#) including web-based trainings, videos, checklists, and references related to vaccine storage and handling.

These example vaccine labels can be used to organize vaccines within the storage unit. Referenced in the storage and handling toolkit.

- [2020-2021 Influenza Season Vaccine Label Examples](#) 📄 [4 pages] (Sept 2020)
- [Vaccine Labels Examples](#) 📄 [20 pages] (Jan 2021)



Vaccine Storage and Handling Toolkit

Updated with 2020-21 Vaccine Storage and Handling Information
Revised and released November 20, 2020



View, download, and print the 2021 Vaccine Storage and Handling Toolkit.

More

Interim Considerations: Preparing for the Potential Management of Anaphylaxis at COVID-19 Vaccination Sites

- Emergency equipment that should be immediately available
- Routine observation periods following COVID-19 vaccination
- Early recognition of anaphylaxis
- Management of anaphylaxis at a COVID-19 vaccination location
- Considerations for anaphylaxis management in special populations
- Patient counseling
- Reporting anaphylaxis

Locations administering COVID-19 vaccines should adhere to CDC guidance, including screening recipients for contraindications and precautions, having necessary supplies and staff members available to manage anaphylaxis, implementing recommended post vaccination observation periods, and immediately treating suspected anaphylaxis with intramuscular epinephrine injection.

CDC v-safe system

- Give patients a **v-safe** information sheet at the time of vaccination
- Suggested healthcare provider script: *CDC has created a way for you to report how you feel after COVID-19 vaccination through a smartphone-based tool that uses text messaging and web surveys to check in with you. Here (or in your packet) is a v-safe information sheet with more details and simple instructions to sign up.*
- Parents and guardians can now enroll adolescents (ages 5 and older) in the v-safe after vaccination health checkerSM and complete health check-ins on their behalf after COVID-19 vaccination.
- Poster and website translated into multiple languages



**Get vaccinated.
Get your smartphone.
Get started with v-safe.**



v-safeSM
after vaccination
health checker

What is v-safe?

v-safe is a smartphone-based tool that uses text messaging and web surveys to provide personalized health check-ins after you receive a COVID-19 vaccination. Through **v-safe**, you can quickly tell CDC if you have any side effects after getting the COVID-19 vaccine. Depending on your answers, someone from CDC may call to check on you. And **v-safe** will remind you to get your second COVID-19 vaccine dose if you need one.

Your participation in CDC's **v-safe** makes a difference—it helps keep COVID-19 vaccines safe.

How can I participate?

Once you get a COVID-19 vaccine, you can enroll in **v-safe** using your smartphone. Participation is voluntary and you can opt out at any time. You will receive text messages from **v-safe** around 2pm local time. To opt out, simply text "STOP" when **v-safe** sends you a text message. You can also start **v-safe** again by texting "START."

How long do v-safe check-ins last?

During the first week after you get your vaccine, **v-safe** will send you a text message each day to ask how you are doing. Then you will get check-in messages once a week for up to 5 weeks. The questions **v-safe** asks should take less than 5 minutes to answer. If you need a second dose of vaccine, **v-safe** will provide a new 6-week check-in process so you can share your second-dose vaccine experience as well. You'll also receive check-ins 3, 6, and 12 months after your final dose of vaccine.

Is my health information safe?

Yes. Your personal information in **v-safe** is protected so that it stays confidential and private.*

Use your smartphone to tell CDC about any side effects after getting the COVID-19 vaccine. You'll also get reminders if you need a second vaccine dose.



Sign up with your
smartphone's browser at

OR

Aim your smartphone's
camera at this code

*To the extent **v-safe** uses existing information systems managed by CDC, FDA, and other federal agencies, the systems employ strict security measures appropriate for the data's level of sensitivity. These measures comply, where applicable, with the following federal laws, including the Privacy Act of 1974; standards enacted that are consistent with the Health Insurance Portability and Accountability Act of 1996 (HIPAA); the Federal Information Security Management Act, and the Freedom of Information Act.

12/01/20

VAERS Reporting and Safety Info - vaers.hhs.gov

VAERS accepts all reports from everyone regardless of the plausibility of the vaccine causing the event or the clinical seriousness of the event.



VAERS Vaccine Adverse Event Reporting System
www.vaers.hhs.gov

About VAERS | Report an Adverse Event | VAERS Data | Resources | Submit Follow-Up Information

Have you had a reaction following a vaccination?

1. Contact your healthcare provider.
2. Report an Adverse Event using the VAERS online form or the downloadable PDF. **New!**

Important: If you are experiencing a medical emergency, seek immediate assistance from a healthcare provider or call 9-1-1. CDC and FDA do not provide individual medical treatment, advice, or diagnosis. If you need individual medical or health care advice, consult a qualified healthcare provider.

¿Ha tenido una reacción después de recibir una vacuna?

1. Contacte a su proveedor de salud.
2. Reporte un evento adverso utilizando el formulario de VAERS en línea. **en Español, Nuevo!**

COVID-19 vaccine EUA reporting requirements for Providers

REPORT AN ADVERSE EVENT
Review reporting requirements and submit reports.

SEARCH VAERS DATA
Download VAERS Data and search the CDC WONDER database.

REVIEW RESOURCES
Find materials, publications, learning tools, and other resources.

SUBMIT FOLLOW-UP INFORMATION
Upload additional information related to VAERS reports.

EUA Fact Sheets

Routine vaccines are given with a Vaccine Information Statement (VIS)

COVID-19 vaccines under the Emergency Use Authorization (EUA) will be given with an EUA Factsheet.

- [Pfizer Bivalent Booster Gray Cap \(Ages 12+, No Diluent\) EUA Fact Sheet](#)
- [Comirnaty/Pfizer Gray Cap \(ages 12+\) EUA Fact Sheet](#)
- [Pfizer Orange Cap \(ages 5-11yrs\) EUA Fact Sheet](#)
- [Pediatric Pfizer Bivalent Booster Orange Cap \(Ages 5 - 11yrs\) EUA Fact Sheet](#)
- [Pfizer Maroon Cap Monovalent & Bivalent \(ages 6mos-4yrs\) EUA Fact Sheet](#)
 - [Pfizer EUA fact sheets translated in other languages](#)
- [Moderna Bivalent Booster Blue Cap \(Ages 6+\) EUA Fact Sheet](#)
- [Moderna Bivalent Booster Dark Pink Cap \(Ages 6mos-5yrs\) EUA Fact Sheet](#)
- [Moderna \(ages 12+\) EUA Fact Sheet](#)
- [Moderna \(ages 6-11\) EUA Fact Sheet](#)
- [Moderna \(ages 6mos-5yrs\) EUA Fact Sheet](#)
 - [Moderna EUA fact sheets translated in other languages](#)
- [Janssen EUA Fact Sheet](#)
 - [Janssen EUA fact sheet translated in other languages](#)
- [Novavax EUA Fact Sheet](#)
 - [Novavax EUA fact sheet translated in other languages](#)

FDA gives Moderna COVID-19 vaccine full approval for use in ages 18 and older

- The Food and Drug Administration (FDA) has given full approval to the Moderna COVID-19 vaccine for use in those ages 18 and older. Ages 6mos to 17 are eligible for Moderna COVID-19 vaccines under FDA emergency use authorization.
- Moderna joins the Pfizer-BioNTech vaccine, which the FDA gave full approval to last August for use in ages 16 and older. Ages 6mos to 15 are eligible for Pfizer COVID-19 vaccines under FDA emergency use authorization.

Moderna Label Update

FDA recently approved updates to both label and the EUA fact sheet for the Moderna COVID-19 vaccine. Distribution of the Moderna vials with the updated labels (see new label above) began during the week of January 13, 2021.

What has changed on the label of the Moderna COVID-19 vaccine:

- The label has been updated to indicate a volume of 5.5mL.
- Doses for primary series and booster shots are now listed on the label.
- Important reminder not to exceed a maximum of 20 punctures per vial

What remains the same for the Moderna COVID-19 vaccine:

- Formulation
- NDC
- Physical vial size
- Number of reportable doses
- Dose volumes (See Below)
- Reporting requirements
 - Wastage should continue to be reported based on 10 doses per vial.
- Proper dosage of Moderna COVID-19 Vaccine
 - Primary series doses 1 and 2 dosage = 0.5 mL
 - Additional primary dose (3rd) dosage for immunocompromised persons = 0.5 mL
 - Booster dosage = 0.25 mL
- Each vial can **ONLY** be punctured up to 20 times.
 - Once the vial has reached the 20-puncture limit, the vial should be discarded **even if there is vaccine remaining in the vial.**



For more information, please refer to the [EUA fact sheet](#) or [Administration Overview for Moderna COVID-19 Vaccine | CDC](#).

Moderna Booster Reporting of Administered and Wasted Doses

- Quick rule of thumb: As long as the vial has been punctured 10 or 14 times and 10 or 14 doses were administered (regardless if they were full primary series doses or half booster doses) there will be no reported wastage.
- Example from the 14 dose vial table: 5 full doses (primary series) were administered and 5 half doses (booster doses) were administered before the 12 hour limit was reached. 4 doses should now be reported as wastage.

Moderna 10 Dose Vial Wastage Table

Full doses \ Half Doses	0	1	2	3	4	5	6	7	8	9	10
0	10	9	8	7	6	5	4	3	2	1	0
1	9	8	7	6	5	4	3	2	1	0	
2	8	7	6	5	4	3	2	1	0		
3	7	6	5	4	3	2	1	0			
4	6	5	4	3	2	1	0				
5	5	4	3	2	1	0					
6	4	3	2	1	0						
7	3	2	1	0							
8	2	1	0								
9	1	0									
10	0										
11											
12											
13											
14											
15											
16											
17											
18											
19											
20											

Green Area = No Waste

Moderna 14 Dose Vial Wastage Table

Full doses \ Half Doses	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
0	14	13	12	11	10	9	8	7	6	5	4	3	2	1	0
1	13	12	11	10	9	8	7	6	5	4	3	2	1	0	
2	12	11	10	9	8	7	6	5	4	3	2	1	0		
3	11	10	9	8	7	6	5	4	3	2	1	0			
4	10	9	8	7	6	5	4	3	2	1	0				
5	9	8	7	6	5	4	3	2	1	0					
6	8	7	6	5	4	3	2	1	0						
7	7	6	5	4	3	2	1	0							
8	6	5	4	3	2	1	0								
9	5	4	3	2	1	0									
10	4	3	2	1	0										
11	3	2	1	0											
12	2	1	0												
13	1	0													
14	0														
15															
16															
17															
18															
19															
20															

No Waste Reported

Green Area = No Waste Reported

Moderna Puncture Tracking Log

- Keep track of how many times vials have been punctured
- Vials can only be punctured maximum of 20 times
- Mark half-doses vs full doses
- Can track 2 vials per log

Moderna Puncture Tracking Log

• • • • •

Vial Lot #: _____

Start Time: _____

End Time: _____

Dose	Full (0.5 mL)	Half (0.25 mL)
1	<input type="checkbox"/>	<input type="checkbox"/>
2	<input type="checkbox"/>	<input type="checkbox"/>
3	<input type="checkbox"/>	<input type="checkbox"/>
4	<input type="checkbox"/>	<input type="checkbox"/>
5	<input type="checkbox"/>	<input type="checkbox"/>
6	<input type="checkbox"/>	<input type="checkbox"/>
7	<input type="checkbox"/>	<input type="checkbox"/>
8	<input type="checkbox"/>	<input type="checkbox"/>
9	<input type="checkbox"/>	<input type="checkbox"/>
10	<input type="checkbox"/>	<input type="checkbox"/>
11	<input type="checkbox"/>	<input type="checkbox"/>
12	<input type="checkbox"/>	<input type="checkbox"/>
13	<input type="checkbox"/>	<input type="checkbox"/>
14	<input type="checkbox"/>	<input type="checkbox"/>
15	<input type="checkbox"/>	<input type="checkbox"/>
16	<input type="checkbox"/>	<input type="checkbox"/>
17	<input type="checkbox"/>	<input type="checkbox"/>
18	<input type="checkbox"/>	<input type="checkbox"/>
19	<input type="checkbox"/>	<input type="checkbox"/>
20	<input type="checkbox"/>	<input type="checkbox"/>

Vial Lot #: _____

Start Time: _____

End Time: _____

Dose	Full (0.5 mL)	Half (0.25 mL)
1	<input type="checkbox"/>	<input type="checkbox"/>
2	<input type="checkbox"/>	<input type="checkbox"/>
3	<input type="checkbox"/>	<input type="checkbox"/>
4	<input type="checkbox"/>	<input type="checkbox"/>
5	<input type="checkbox"/>	<input type="checkbox"/>
6	<input type="checkbox"/>	<input type="checkbox"/>
7	<input type="checkbox"/>	<input type="checkbox"/>
8	<input type="checkbox"/>	<input type="checkbox"/>
9	<input type="checkbox"/>	<input type="checkbox"/>
10	<input type="checkbox"/>	<input type="checkbox"/>
11	<input type="checkbox"/>	<input type="checkbox"/>
12	<input type="checkbox"/>	<input type="checkbox"/>
13	<input type="checkbox"/>	<input type="checkbox"/>
14	<input type="checkbox"/>	<input type="checkbox"/>
15	<input type="checkbox"/>	<input type="checkbox"/>
16	<input type="checkbox"/>	<input type="checkbox"/>
17	<input type="checkbox"/>	<input type="checkbox"/>
18	<input type="checkbox"/>	<input type="checkbox"/>
19	<input type="checkbox"/>	<input type="checkbox"/>
20	<input type="checkbox"/>	<input type="checkbox"/>

Reminders

- Moderna booster doses are 0.25mL (half the dose size of a primary series dose)
- Moderna vials can only be punctured a maximum of 20 times
- Providers are responsible to keep track of the number of punctures for each vial
- Punctured Moderna vials can be used to administer both primary and booster doses
- Punctured Moderna vials must be used within 12 hours

602-364-3899

ArizonaVFC@azdhs.gov

azdhs.gov/covid19/vaccines

Updated 12/2021

You can only puncture the diluent once

Orders of Pfizer age 5-11 vaccine include ancillary kits that contain 10mL diluent vials. While these vials appear to contain sufficient diluent for multiple vials, they must only be used once.

- Diluent vials are a one-time-use item and should be discarded with the remaining content after each use.
- For each vial of vaccine, extract 1.3mL of diluent from a single-use vial to reconstitute 1.3mL of vaccine
- Do not be tempted to puncture diluent vials more than once.

COVID-19 Vaccine Transfer Matchmaker Tool

- **Share vaccine with other active COVID-19 providers**
- Request vaccine from nearby locations if active providers need vaccine
- You can transfer needles as well
- Only serves as connection between providers
- **All transfers must still have prior approval in ASIIS, must follow all transfer steps**
- **Matchmaker Tool Job Aid**
 - Transfer steps
 - Helpful resources linked

COVID-19 Vaccine Transfer Matchmaker



Do you have more vaccines than you can use?

List them on the [Arizona COVID-19 Vaccine Transfer Matchmaker Website](#) using the "Add Vaccine/Needle" form on the website.



Do you need to place a vaccine order?

First, check the [Arizona COVID-19 Vaccine Transfer Matchmaker Website](#) for vaccine available at a location near you, and submit a "Request Vaccine/Needle" form on the website. The table on the left shows vaccines that are available for providers to request.

72,182
Vaccines requested to date

How ID	County	City	Vaccine/Needle	Expiration Date	Quantity
1420	Maricopa	Phoenix	Pfizer age 12+	11/03/21	127886
1423	Maricopa	Phoenix	Pfizer age 12+	11/03/21	13462
1424	Maricopa	Phoenix	Pfizer age 12+	11/03/21	10302
1419	Maricopa	Scottsdale	J&J	03/03/22	80
1416	Maricopa	Phoenix	Pfizer age 12+	11/03/21	81154
1416	Maricopa	Phoenix	Pfizer age 12+	11/03/21	84300
1389	Maricopa	Phoenix	J&J	04/09/22	50
1389	Cochise	Flagstaff	Pfizer age 12+	11/03/21	3010
1388	Pima	Green Valley	Pfizer age 12+	11/03/21	870
1372	Maricopa	Phoenix	Pfizer age 12+	11/03/21	36124
1004	Yuma	Yuma	Pfizer age 12+	12/31/21	1080

Add Vaccine/Needle

What is your name? *

What is your email address? *

What is your phone number? *

What is your ASIS PIN? *

What city are you located in? *

What county are you located in? *

Which vaccine/needle is available? *

Auxiliary kits must be transferred with the vaccine. Please do NOT list needles separately if you are already listing vaccines.

☐ J&J ☐ Moderna ☐ Pfizer age 12+ ☐ Pfizer age 11-1 ☐ 1* needles only (no vaccines) ☐ 1.5* needles only (no vaccines)

How many doses/needles? *

Request Vaccine/Needle

What is your name? *

What is your email address? *

What is your phone number? *

What is your ASIS PIN? *

What is the row ID that you are interested in? *

Auxiliary kits must be transferred with the vaccine. You do NOT need to request needles separately if you are requesting vaccines.

How many doses/needles? *

☐ Send me a copy of my responses

[Privacy Notice](#) | [Report Abuse](#)

ADHS Consent Form Template

- No Federal requirement for informed consent relating to immunization
- Available for providers who do not have own consent form
- Use with [CDC Pre-Vaccination checklist](#)
- [Spanish ADHS consent form](#)
- [Spanish CDC Pre-Vaccination Checklist](#)

Prevaccination Checklist for COVID-19 Vaccines

For vaccine recipients: The following questions will help us determine if there is any reason you should not get the COVID-19 vaccine today. **If you answer "yes" to any question, it does not necessarily mean you should not be vaccinated.** It just means additional questions may be asked. If a question is not clear, please ask your healthcare provider to explain it.

Patient Name _____ Age _____

	Yes	No	Don't know
1. Are you feeling sick today?			
2. Have you ever received a dose of COVID-19 vaccine?			
• If yes, which vaccine product did you receive? <input type="checkbox"/> Pfizer <input type="checkbox"/> Moderna <input type="checkbox"/> Janssen (Johnson & Johnson) <input type="checkbox"/> Another product _____			
3. Have you ever had an allergic reaction to: (This would include a severe allergic reaction [e.g., anaphylaxis] that required treatment with epinephrine or EpiPen® or that caused you to go to the hospital. It would also include an allergic reaction that occurred within 4 hours that caused hives, swelling, or respiratory distress, including wheezing.)			

ADHS COVID-19 Vaccine Consent Form



Use this form in conjunction with the [CDC Pre-Vaccination Checklist for COVID-19 Vaccines](#).

(Staff only) Appointment ID: _____

Patient Information

Last Name		First Name		Middle Name (optional)
Mother's Maiden Name (Optional)		Date of Birth (MM/DD/YYYY)		Gender
Address		Apartment Number	City	State Zip
<input type="radio"/> No address available		Phone Number		

Insurance Information

Do you have insurance? ☐ Yes ☐ No

Plan Name Plan Group ID # Plan Individual ID #

Name of Person Covered By Plan Plan Responsible Person Name

Private Insurance Address and Phone Number (If Available)

CONSENT AND ASSIGNMENT OF BENEFITS: I have had a copy of the Emergency Use Authorization for the COVID-19 vaccine made available to me. I have had a chance to ask questions and I believe I understand the benefits and risks of the COVID-19 vaccines requested. I ask that the vaccines be administered to me or the person for whom I am authorized to make this request.

I certify that I am: (1) the patient and at least 18 years of age; (2) the legal guardian of the patient and the patient's age makes him/her eligible to receive the vaccine based on the current emergency use authorization; or (3) a person authorized to consent on behalf of the patient where the patient is unable to consent for themselves.

I hereby assign to _____ any insurance or other third-party benefits available for the administration fee of the COVID-19 vaccine provided to me. I agree to forward to _____ all health insurance and other third-party payments I receive for services rendered to me immediately upon receipt.

I agree to allow the health care provider to release information to the Arizona State Immunization Information System (ASIS) to record that I (or for the person for whom I am authorized to consent) have received this COVID-19 vaccine. This information will help keep track of the manufacturer and doses of the vaccine.

Patient Printed Name	Patient Signature	Date Signed
Parent/Guardian/Authorized Person Printed Name	Authorized Person's Signature	Date Signed

Vaccine Administration Information for Immunizer Use Only


Administration Date	Manufacturer	NDC #	<input type="radio"/> LEFT ARM <input type="radio"/> RIGHT ARM
Lot Number	Expiration Date	Route	Site
Administering Immunizer Name and Title		Administering Immunizer Signature	
Is this the patient's first, second, or third dose? <input type="radio"/> First <input type="radio"/> Second <input type="radio"/> Third <input type="radio"/> Booster Dose			

ADHS Does Not Provide Exemption Forms for Adults for the COVID-19 Vaccine


- Individual facilities and employers may choose to set their own immunization requirements for adults
- If an employer is requiring a COVID-19 Vaccine for their employees, and the employee is refusing vaccination, direct that patient to their Human Resources department
- Adults who have questions on immunization exemptions, for any vaccine, should contact the facility or employer who has set the requirement

Standing Order Links - Pfizer-BioNTech

Link to Pfizer (Gray Cap)



12 Years of Age and Older (Gray Cap)
Pfizer-BioNTech COVID-19 Vaccine
Standing Orders for Administering Vaccine



Vaccine	Diluent	Dosage (amount)/ Route
12 years of age and older (gray cap)	No NOT dilute.	0.2 mL/IM injection

Purpose

- To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

Policy


- Where authorized under state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess and vaccinate persons who meet the criteria in the "Procedure" section below without the need for clinician examination or direct order from the attending provider at the time of the interaction.

Procedure


Assess persons 12 years of age and older for vaccination with Pfizer-BioNTech COVID-19 Vaccine based on the following criteria:

- Persons who **ARE NOT** moderately or severely immunocompromised*
- If the recipient has never received a COVID-19 vaccine, administer 1 dose of Pfizer-BioNTech COVID-19 Vaccine.
- If the recipient has received 1 previous dose of:
 - Pfizer-BioNTech COVID-19 Vaccine, administer the second dose at least 21 days (3 weeks) after the first dose.
 - If the first-dose vaccine product cannot be determined or is no longer available, administer Pfizer-BioNTech COVID-19 Vaccine at least 21 days (3 weeks) after the first dose.
- If the recipient has received:
 - 2 doses of Pfizer-BioNTech Vaccine, administer a third dose at least 28 days (4 weeks) after dose 2.
 - 1 dose of Janssen COVID-19 Vaccine, administer Pfizer-BioNTech COVID-19 Vaccine at least 28 days (4 weeks) after Janssen COVID-19 Vaccine.
- If the recipient has received:
 - 3 previous doses of Pfizer-BioNTech COVID-19 Vaccine, administer a booster dose at least 3 months after the dose 3 in the mRNA primary series.
 - 1 dose of Janssen and an additional dose of mRNA vaccine (any product), administer a booster dose at least 2 months (8 weeks) after the additional dose of mRNA vaccine.
- If the recipient has received 2 previous doses of Pfizer-BioNTech COVID-19 Vaccine, administer a booster dose at least 3 months after the second dose.

Link to Pfizer (Orange Cap)



5 Through 11 Years of Age
Pfizer-BioNTech COVID-19 Vaccine
Standing Orders for Administering Vaccine



Vaccine	Diluent	Dosage (amount)/ Route
5 through 11 years of age	1.3 mL of 0.9% sodium chloride (normal saline, preservative-free) diluent	0.2 mL/IM injection

Purpose

- To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

Policy

- Where authorized under state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess and vaccinate persons who meet the criteria in the "Procedure" section below without the need for clinician examination or direct order from the attending provider at the time of the interaction.

Procedure

Assess children 5 through 11 years of age for vaccination with Pfizer-BioNTech COVID-19 Vaccine based on the following criteria:


- Primary series***
 - If the recipient has never received a COVID-19 vaccine, administer 1 dose of Pfizer-BioNTech COVID-19 Vaccine.
- Booster dose***
 - If the recipient has received 1 previous dose of Pfizer-BioNTech COVID-19 Vaccine, administer the second dose at an interval of at least 21 days.
- Children who are **NOT** immunocompromised: If the recipient has received 2 previous doses of Pfizer-BioNTech COVID-19 Vaccine, administer a booster dose at least 5 months after dose 2.
- Children who are **moderately or severely immunocompromised**†: If the recipient has received 2 previous doses of Pfizer-BioNTech COVID-19 Vaccine, administer a booster dose at least 3 months after dose 2.
- Children with a history of myocarditis or pericarditis:
 - If history is prior to COVID-19 vaccination, may receive Pfizer-BioNTech from 12 months to 14 weeks of age.

Screen for contraindications and precautions


Contraindications:

- History of:
 - Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to component of the COVID-19 vaccine.
 - Known diagnosed allergy to a component of the vaccine.

Link to Pfizer (Maroon Cap)



6 Months Through 4 Years of Age
Pfizer-BioNTech COVID-19 Vaccine
Standing Orders for Administering Vaccine



Vaccine	Diluent	Dosage (amount)/ Route
6 months through 4 years of age	2.2 mL of 0.9% sodium chloride (normal saline, preservative-free) diluent	0.2 mL/IM injection

Purpose

- To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

Policy

- Where authorized under state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess and vaccinate persons who meet the criteria in the "Procedure" section below without the need for clinician examination or direct order from the attending provider at the time of the interaction.

Procedure

Assess children 6 months through 4 years of age for vaccination with Pfizer-BioNTech COVID-19 Vaccine based on the following criteria:

- Children who **ARE NOT** moderately or severely immunocompromised*
- Primary series***
 - If the recipient has never received a COVID-19 vaccine, administer 1 dose of Pfizer-BioNTech COVID-19 Vaccine.
- If the recipient has received 1 previous dose of Pfizer-BioNTech COVID-19 Vaccine, administer the second dose at least 3-8 weeks after Dose 1.
- If the recipient has received 2 previous doses of Pfizer-BioNTech Vaccine, administer the third dose at least 8 weeks after Dose 2.

Screen for contraindications and precautions

Contraindications:

- History of:
 - Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to component of the COVID-19 vaccine.
 - Known diagnosed allergy to a component of the vaccine.

[Link to Moderna](#)
[Ages 6mos - 5yrs](#)

6 Months Through 5 Years of Age

Moderna COVID-19 Vaccine

Standing Orders for Administering Vaccine

Vaccine	Dosage (amount) / Route
Formulation: 6 months through 5 years of age	0.25 mL (mL) / Intramuscular

Purpose

To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

Policy

Where authorized under state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess and vaccinate persons who meet the criteria in the "Indications" section below without the need for clinical examination or direct order from the attending provider at the time of the vaccination.

Precautions

Issues children 6 months through 5 years of age for vaccination with Moderna COVID-19 vaccine based on the following criteria:

- Children who are NOT moderate or severely immunocompromised¹
 - Primary Series
 - If the recipient has never received a COVID-19 vaccine, administer 1 dose of Moderna COVID-19 Vaccine.
 - If the recipient has received 1 previous dose of Moderna COVID-19 vaccine, administer the second dose at least 4–8 weeks after Dose 1.
- Children who ARE moderate or severely immunocompromised²
 - Primary Series
 - If the recipient has never received a COVID-19 vaccine, administer 1 dose of Moderna COVID-19 Vaccine.
 - If the recipient has received 1 previous dose of Moderna COVID-19 vaccine, administer the second dose at least 4

- If mycoplasmas or pericarditis occurred after the first dose of mRNA vaccine, discuss advice on additional doses of Moderna COVID-19 vaccine. If after a risk assessment, a decision is made to administer a subsequent dose of Moderna COVID-19 vaccine, vaccine should not be administered until the mycoplasmas or pericarditis has resolved. Clinical considerations can be found at <https://www.cdc.gov/covid-19/clinical-considerations/after-biotech-moderna>.

Additional clinical considerations

- For children who received a COVID-19 vaccine that is not currently authorized or approved in the United States, guidance can be found at: <https://www.cdc.gov/covid-19/clinical-considerations/18mths-to-19mths-not-currently-authorized-vaccines.html#first-authorized-vaccines>.
- Moderna COVID-19 Vaccine may be administered with other vaccines without the need to timing including simultaneous administration.
- For recommendations for COVID-19 vaccination and SARS-CoV-2 infection, visit <https://www.cdc.gov/covid-19/clinical-considerations/covid-19-infection-and-vaccination.html#coVid-19-clinical-considerations>

Screen for contraindications and precautions

Contraindications:

- History of:
 - Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a component of the COVID-19 vaccine
 - Known diagnosed allergy to a component of the vaccine
- For more information on contraindications, clinical considerations, interim considerations, see <https://www.cdc.gov/covid-19/clinical-considerations/interim-considerations.html#appendix>

Precautions:

Standing Order Links - Janssen

[Link to Janssen](#)

Vaccine	Dosage (amount)/ Route
Janssen COVID-19 Vaccine (Johnson & Johnson)	0.5 mL/IM injection

Janssen COVID-19 Vaccine
(Johnson & Johnson)
Standing Orders for Administering Vaccine
to Persons 18 Years of Age and Older

Purpose

- To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

Policy

- Where authorized under state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess and vaccinate persons who meet the criteria in the "Procedure" section below without the need for clinician examination or direct order from the attending provider at the time of the interaction.

Procedure

Assess persons 18 years of age and older for vaccination with Janssen COVID-19 Vaccine based on the following criteria:

- If the recipient has never received a COVID-19 vaccine, administer 1 dose of Janssen COVID-19 vaccine.
- If the recipient has received 1 dose of a Janssen COVID-19 Vaccine, no additional primary-series doses are needed. A booster dose is recommended 2 months (8 weeks) after the primary Janssen dose; any FDA-authorized or approved COVID-19 vaccine may be given.
- If 2 doses of an mRNA vaccine or a single dose of Janssen COVID-19 Vaccine has been administered, the person is considered fully vaccinated 2 weeks after completing the primary vaccination series.
- In situations where the first dose of an mRNA COVID-19 vaccine was received but the patient is unable to complete the series with either the same or different mRNA COVID-19 vaccine, (e.g., due to contraindication) consideration may be given to vaccination with the Janssen COVID-19 Vaccine at a minimum interval of 28 days after receipt of mRNA COVID-19 vaccine dose. However, vaccination should be done in an appropriate setting under the supervision of a healthcare provider experienced in the management of severe allergic reactions. Consider referral to an allergist-immunologist.

Thrombocytopenia syndrome (TTS) and thrombocytopenia:


- Inform women aged 18-49 years of the increased risk of thrombosis with thrombocytopenia syndrome (TTS) in their age group after Janssen COVID-19 vaccination and about the availability of other authorized vaccines (i.e., mRNA vaccines).
- A second dose of Janssen COVID-19 vaccine is not recommended for people who had TTS after their first dose. These people may receive a dose of mRNA COVID-19 vaccine as a booster dose at least 2 months (8 weeks) after their dose of Janssen vaccine and after their clinical condition has stabilized.¹
- A consultation with the patient's clinical team, including hematologists or other specialists, should be considered.
- Offer another FDA-authorized or approved vaccine (i.e., mRNA vaccine) to unvaccinated persons with a history of an episode of an immune-mediated syndrome characterized by thrombosis and thrombocytopenia (e.g., heparin-induced thrombocytopenia) if it has been 90 days or less since their illness resolved. After 90 days, patients may be vaccinated with any FDA-authorized or approved COVID-19 vaccine.

NOTE: Persons at risk or with a history of other thrombosis not associated with thrombocytopenia can receive an FDA-authorized or approved vaccine

- People with a history of Guillain-Barré Syndrome (GBS):
- Can receive any FDA-authorized or approved COVID-19 vaccine. However, given the possible association between the Janssen COVID-19 vaccine and an increased risk of GBS, discuss with these patients the availability of mRNA COVID-19 vaccines that offer protection against COVID-19.
- Persons who have received HCT or CAR-T-cell therapy
- Revacinate persons who received doses of COVID-19 vaccine prior to receiving HCT or CAR-T-cell therapy with a primary series at least 3 months (12 weeks) after transplant or CAR-T-cell therapy.
- Booster doses:
- Administer a booster dose at least 2 months (8 weeks) after completion of the Janssen COVID-19 Vaccine primary dose to:


Standing Order Links - Novavax

Link to Novavax



Novavax COVID-19 Vaccine

Standing Orders for Administering Vaccine
to Persons 18 Years of Age and Older



Vaccine Product	Dose/Injection Amount	Route
Blue capped vial	5 µg I5 and 50 µg of Matrix-M™ adjuvant/0.5 mL	Intramuscular (IM) injection

Purpose

- To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

Policy

- Where authorized under state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess and vaccinate persons who meet the criteria in the "Procedure" section below without the need for clinician examination or direct order from the attending provider at the time of the interaction.

Procedure

Assess persons 18 years of age and older for vaccination with Novavax COVID-19 Vaccine based on the following criteria:

Persons who are **NOT** moderately or severely immunocompromised¹

- If the recipient has never received a COVID-19 vaccine, administer 1 dose of Novavax COVID-19 Vaccine.
- If the recipient has received 1 previous dose of:
 - Novavax COVID-19 Vaccine, administer the second dose at least 3 to 8 weeks after the first dose.
 - A vaccine product that cannot be determined, is no longer available or contraindicated, administer Novavax COVID-19 Vaccine at least 4-8 weeks after the first dose.

Persons who are **ARE** moderately or severely immunocompromised¹

- If the recipient has never received a COVID-19 vaccine, administer 1 dose of Novavax COVID-19 Vaccine.
- If the recipient has received 1 previous dose of:
 - Novavax COVID-19 Vaccine, administer the second dose at least 3 weeks after Dose 1.
 - A vaccine product that cannot be determined, is no longer available or contraindicated, administer Novavax COVID-19 Vaccine at least 4 weeks after the first dose.

- Persons with a history of myocarditis or pericarditis:
 - If history is prior to COVID-19 vaccination, may receive Novavax COVID-19 vaccine, after the episode of myocarditis or pericarditis has completely resolved.
 - If myocarditis or pericarditis occurred after the first dose of an mRNA or Novavax COVID-19 vaccine, generally experts advise no additional doses of any COVID-19 vaccine. Administration of the second dose of an mRNA or Novavax COVID-19 vaccine series can be considered in certain circumstances after the episode of myocarditis or pericarditis has completely resolved. Considerations can be found at <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#myocarditis-pericarditis>
- Additional clinical considerations
 - For people who received a COVID-19 vaccine outside the United States, guidance can be found at: <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a>
 - Novavax COVID-19 Vaccine may be coadministered with other routinely recommended vaccines without regard to timing, including simultaneous administration.
 - For recommendations for COVID-19 vaccination and SARS-CoV-2 infection, see <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html#infection>
- Persons who have received HCT or CAR-T-cell therapy
 - Revaccinate persons who received doses of COVID-19 vaccine prior to or during receiving HCT or CAR-T-cell therapy with a primary series at least 3 months (12 weeks) after transplant or CAR-T-cell therapy.
- Screen for contraindications and precautions.
 - Contraindications**
 - History of a:
 - Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a component of the COVID-19 vaccine
 - Known diagnosed allergy to a component of the vaccine (see <https://www.cdc.gov/vaccines/covid-19/clinical-considerations-us-appendix.html#appendix-a>)

ADHS Vaccine Reference Sheets - Janssen

Janssen (Ages 18+)

- [Janssen Vaccine Reference Sheet](#)
- [What Patients and Providers Need to Know About the Janssen Vaccine](#)

Vaccine Storage

Janssen COVID-19 Vaccine

Refrigerator



- 2.0° C to 8.0° C
- 36.0° F to 46.0° F
- Store in refrigerator for up to 11 months
- DO NOT REFREEZE

Dosage



- Each vaccine vial contains 5, 0.5mL doses to be administered
- No reconstitution required
- Discard any punctured vial held at refrigerator temperatures after 6 hours
- Discard any punctured vial held at room temperature (maximally 25° C / 77° F) after 2 hours

Administration




- Standard needles and syringes
- Draw 0.5mL and inject intramuscularly (IM)
- Single dose
- For adults 18 and older
- Not for children
- Can be administered until 11:59 p.m. EST on expiry date. Expiry dates can be verified on the [Janssen website](#)

The storage and handling information found on the Fact Sheet supersedes the storage and handling information on the carton and vial labels.

Vaccine Manufacturer COVID-19 Websites


- [Comirnaty/Pfizer COVID-19 Website](#)
- [Moderna COVID-19 Website](#)
- [Janssen COVID-19 Website](#)
- [Novavax COVID-19 Website](#)

Pfizer-BioNTech COVID-19 Vaccine  English

Global Information About Pfizer-BioNTech COVID-19 Vaccine (also known as BNT162b2)

The approval status of the Pfizer-BioNTech COVID-19 Vaccine varies worldwide. In countries where the vaccine has not been approved by the relevant regulatory authority, it is an investigational drug, and its safety and efficacy have not been established.


As country information may vary, please choose the country below in which you are a licensed healthcare professional for more information on the Pfizer-BioNTech COVID-19 Vaccine.



This site is intended for Healthcare Professionals only.

I am a Healthcare Professional in:

Select



I am NOT a Healthcare Professional

Select

[Report an Adverse Event](#)

Information about the Pfizer-BioNTech COVID-19 Vaccine is only available for certain countries. This site will be updated as more information becomes available.

Moderna COVID-19 Vaccine

moderna

GO TO FULL UNITED STATES SITE HERE



Novavax COVID-19 Vaccine (recombinant, adjuvanted)

For US Vaccination Providers ▶

For US Vaccine Recipients ▶

Global information about the Novavax COVID-19 Vaccine (recombinant, adjuvanted)


The approval status of the Novavax COVID-19 Vaccine (recombinant, adjuvanted) (also known as NVX-CoV2373) varies worldwide. In countries where the vaccine has not been approved by the relevant regulatory authority or regulatory process, you will not find information about it on this site. This site will be updated as more information becomes available.

- The Moderna COVID-19 Vaccine has not been approved or licensed by the U.S. Food and Drug Administration (FDA), but has been authorized for emergency use by FDA, under an Emergency Use Authorization (EUA), to prevent Coronavirus Disease 2019 (COVID-19) for individuals 18 years of age and older. There is no FDA-approved vaccine to prevent COVID-19.
- The EUA for the Moderna COVID-19 Vaccine is in effect for the duration of the declaration justifying emergency use of the product, unless the declaration is terminated or authorization is revoked sooner.

Download EUA Fact Sheet & Full PI for Vaccination Providers

Download EUA Fact Sheet for Vaccine Recipients & Caregivers

Look Up Vaccine Expiration Dates for Vaccination Providers

janssen  Important Safety Information [Janssen.com](#) [Contact Us](#)

The U.S. FDA Has Granted the Janssen COVID-19 Vaccine an Emergency Use Authorization (EUA)

To learn more, choose the appropriate option:

I am a healthcare provider

I am NOT a healthcare provider

The Janssen COVID-19 Vaccine has not been approved or licensed by the U.S. Food and Drug Administration (FDA), but has been authorized by FDA through an Emergency Use Authorization (EUA) for active immunization to prevent Coronavirus Disease 2019 (COVID-19) in individuals 18 years of age and older. The emergency use of this product is authorized only for the duration of the declaration that circumstances exist justifying the authorization of the emergency use of the medical product under Section 564(b)(1) of the FDCA Act, unless the declaration is terminated or authorization revoked sooner.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

Do not administer the Janssen COVID-19 Vaccine to individuals with a known history of a severe allergic reaction (e.g., anaphylaxis) to any component of the Janssen COVID-19 Vaccine.

WARNINGS AND PRECAUTIONS

- **Management of Acute Allergic Reactions:** Appropriate medical treatment to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of the Janssen COVID-19 Vaccine.

Monitor Janssen COVID-19 vaccine recipients for the occurrence of adverse events and reactions according to the Centers for Disease Control and Prevention guidelines.

CDC COVID-19 Vaccine Resources

- [CDC Pfizer \(Comirnaty\) Resource Page](#)
- [CDC Moderna Resource Page](#)
- [CDC Janssen Resource Page](#)
- [CDC Novavax Resource Page](#)

12 Years of Age and Older Pfizer-BioNTech COVID-19 Vaccine

Vaccine Preparation and Administration Summary

General Information

Vaccine: Pfizer-BioNTech COVID-19 Vaccine (purple cap)

Use the correct formulation based on the age of the recipient

Diluent: 1.8 mL of 0.9% sodium chloride (normal saline, preservative-free) Use a new vial every time.

Multidose vial: 6 doses per vial

Dosage: 0.3 mL

Prepare the vaccine using a NEW vial of diluent EVERY TIME. Discard the diluent vial and remaining diluent after mixing the vaccine.

Age Indications

12 years of age and older

Thawing Frozen Vaccine

- Frozen vaccine must be thawed before using.
- Thaw vaccine in the refrigerator or at room temperature:
 - Refrigerator:** Between 2°C and 8°C (36°F and 46°F) Unpunctured vials may be stored in the refrigerator for up to 1 month (31 days).
 - Room temperature (for immediate use):** Up to 25°C (77°F) Unpunctured vials cannot be kept at room temperature for more than 2 hours (including thaw time).
- Amount of time needed to thaw vaccine varies based on temperature and number of vials.

Prepare the Vaccine

Follow aseptic technique. Perform hand hygiene before vaccine preparation, between patients, when changing gloves (if worn), and any time hands become soiled.*

Remove vaccine from the storage unit. Check the vial label to ensure it is the correct formulation based on the age of the recipient. The vial for persons 12 years of age and older has a purple cap and purple border on the label. Allow vaccine to come to room temperature. Vials can be held at room temperature for up to 2 hours before mixing.

Before mixing, check:

- expiration date on the vaccine and diluent
- any beyond-use dates

NEVER use expired vaccine or diluent. **NEVER** use vaccine after the beyond-use date or times.

With the vaccine at room temperature, gently invert vial 10 times. Do not shake the vial. If the vial is shaken, contact the manufacturer. The vaccine is white to off-white in color and may contain opaque particles. Do not use if liquid is disclosed.

Using a new, sterile alcohol prep pad for each vial, wipe off the stoppers of the diluent and vaccine vials. Using a 21-gauge (or narrower) needle, withdraw 1.8 mL of 0.9% sodium chloride (normal saline, preservative-free) into a mixing syringe. Discard diluent vial and any remaining diluent every time. Do NOT use bacteriostatic normal saline or other diluents to mix the vaccine.

1.8 mL

Janssen COVID-19 Vaccine (Johnson & Johnson)

Vaccine Preparation and Administration Summary

General Information

Vaccine: Janssen COVID-19 Vaccine (Johnson & Johnson)

Administration

Intramuscular (IM) injection in the deltoid muscle

at room temperature: 8°C (36°F and 46°F).

at room temperature for up to 30 minutes

moving vials from the freezer. for this vaccine to track ratures.

an the QR code located ll bring up a website; r the lot number, and the other option is to access the <https://www.cdc.gov/pfizer/downloads/expiration>.

ing expiration dates.

Novavax COVID-19 Vaccine

Vaccine Preparation and Administration Summary

General Information

* Novavax is not authorized for ANY booster doses including any series.

Moderna COVID-19 Vaccine

Vaccine Preparation and Administration Summary

General Information

Vaccine: Moderna COVID-19 Vaccine (Johnson & Johnson)

Administration

Intramuscular (IM) injection in the deltoid muscle

the vaccine vial or carton.

ter carton, or

in date approaches, check ard vaccine until ensuring DC's expiration date date changes.

ore using.

at room temperature: 8°C (36°F and 46°F).

d in the refrigerator

°C and 25°C (46°F and 77°F).

at room temperature for up to 30 minutes

accine varies based on

r and 30 minutes

moving vials from the freezer. for this vaccine to track ratures.

an the QR code located ll bring up a website; r the lot number, and the other option is to access the <https://www.cdc.gov/pfizer/downloads/expiration>.

ing expiration dates.

COVID-19 Vaccine Expiration Dates - Pfizer

Pediatric Pfizer Maroon Cap Monovalent and Bivalent (Ages 6 months - 4 years)

- Date printed on vial and tray is **manufacturing date only**
- Use the [Pfizer website](#)

Base expiration date on storage method and manufacturing date. For example:

- If the manufacturer date is 8/2021 and the doses are stored in an ultra-low-temp freezer, the doses would expire April 30, 2022.
- If the doses are put in the refrigerator November 2, 2021, they would expire 10 weeks from November 2 on January 11, 2022.

Pfizer Orange Cap (Ages 5-11)

- Date printed on vial and tray is **manufacturing date only**
- Use the [Pfizer Age 5-11 EUA Fact Sheet](#) (Page 3)
- Use the [Pfizer website](#)

Base expiration date on storage method and manufacturing date. For example:

- If the manufacturer date is 8/2021 and the doses are stored in an ultra-low-temp freezer, the doses would expire April 30, 2022.
- If the doses are put in the refrigerator November 2, 2021, they would expire 10 weeks from November 2 on January 11, 2022.

Pfizer Gray Cap (Ages 12+)

- Date printed on vial and tray is **manufacturing date only**
- Use the [Pfizer Gray Cap \(Ages 12+\) EUA Fact Sheet](#) (Page 5)
- Use the [Pfizer website](#)

Base expiration date on storage method and manufacturing date. For example:

- If the manufacturer date is 8/2021 and the doses are stored in an ultra-low-temp freezer, the doses would expire April 30, 2022.
- If the doses are put in the refrigerator November 2, 2021, they would expire 10 weeks from November 2 on January 11, 2022.

Comirnaty Gray Cap (Ages 12+)

- Date printed on vial and tray is the **expiration date**
- Use the [Pfizer website](#) to double check expiration dates

Pfizer-BioNTech Bivalent Booster (Ages 12+, No Diluent) - Gray Cap

- Date printed on vial and tray is the **expiration date**
- Use the [Pfizer website](#) to double check expiration dates

Pediatric Pfizer-BioNTech Bivalent Booster (Ages 5 - 11 years) - Orange Cap

- Date printed on vial and tray is the **expiration date**
- Use the [Pfizer website](#) to double check expiration dates

COVID-19 Vaccine Expiration Dates - Moderna, Novavax and Janssen

Pediatric Moderna **Blue Cap** with **Magenta Border** on label (Ages 6 months to 5 years)

- Use QR code
- Enter lot number on [expiration date checker website](#)

Pediatric Moderna **Blue Cap** with **Purple Border** on label (Ages 6 to 11 years)

- Use QR code
- Enter lot number on [expiration date checker website](#)

Moderna **Bivalent Booster Dark Pink Cap** with **Yellow Border** on label (Ages 6 months - 5 years)

- Use QR code
- Enter lot number on [expiration date checker website](#)

Moderna **Bivalent Booster Blue Cap** (Ages 6+)

- Use QR code
- Enter lot number on [expiration date checker website](#)

Moderna **Red Cap** (Ages 12+)

- Use QR code
- Enter lot number on [expiration date checker website](#)

Janssen

- Use QR code
- Enter lot number on [expiration date checker website](#)
- Can be used until 11:59 EST on expiry date

Novavax

- Use QR code
- Enter lot number on [expiration date checker website](#)

CDC Expiration Date Tracking Tool

- Pfizer Gray Cap (ages 12+) , **Orange Cap** (ages 5-11yrs) & **Maroon Cap** (ages 6mo - 4yrs) doses can be refrigerated for up to 10 weeks
- Moderna doses can be refrigerated for up to 30 days
- Janssen doses can be refrigerated for up to 11 months
- Providers should check the latest expiry information on the manufacturer's website before doses are removed from the unit

[illegible]

COVID-19 Vaccine and Beyond-use Dates (BUDs)

- CDC has developed tracking labels refrigerators and freezers to help monitor and document cold storage dates
- Assist with documenting transportation time and temperature
- Available for both Moderna and Pfizer/Comirnaty vaccines
- [Moderna COVID-19 Vaccine: Beyond Use Date/Time \(BUD\) Tracking Label for Vaccine During Refrigerator Storage \(cdc.gov\)](#)
- [Pfizer-BioNTech COVID-19 Beyond Use Date/Time \(BUD\) Tracking Labels for Vaccine During Freezer or Refrigerator Storage \(cdc.gov\)](#)

Returning Thermal Shipping Containers (Once Empty)

Pfizer

- Shippers **DO NOT** need to be returned
- Only controlant data logger needs to be returned

Moderna

- Shippers do not need to be returned

Janssen

- Shippers do not need to be returned

Novavax

- Shippers do not need to be returned

Vaccine Process

Vaccines are widely available

- Providers enter their COVID-19 vaccine order in ASIIS
- BIZS will approve the order
- The order goes to CDC to Distributor/Manufacturer
- The order is shipped (will show in ASIIS)
- Provider receives shipment of ancillary kit and vaccine (may not be same day)
- Provider logs into ASIIS to mark doses as “received”
- Provider administers vaccine
- Provider reports the administered vaccine to ASIIS through their EHR, State VMS POD application, directly in ASIIS or using Mass Immunization Module in ASIIS
- The dose is subtracted from the ASIIS inventory if properly recorded in a patient’s ASIIS record

ASIIS Information

Placing Orders in ASIIS

- Providers who have an approved onboarding submission are able to order vaccines
- AIPO Train courses
 - How to Place an Order in ASIIS
 - Vaccine Inventory Management - information on ordering and inventory reconciliation
 - Sign up for Dose Accountability Webinar
 - Recordings available any time within AIPO Train course
 - Included in Pandemic Provider course bundle
- Use COVID-19 Vaccine Presentations Job Aid to help make ordering decisions

ARIZONA DEPARTMENT OF HEALTH SERVICES

COVID-19 Vaccine Presentations Available to Order in ASIIS

Pfizer

Comirnaty (Ages 12+, No Diluent) - Gray Cap
180-dose minimum order
• 6-dose multi-dose vial
• NDC 00069-2025-10

Pfizer-BioNTech Bivalent Booster
180-dose minimum order
• 6-dose multi-dose vial
• NDC 59267-0304-02

Pfizer-BioNTech Bivalent Booster Dose Vial
50-dose minimum order
• Single dose vial
• NDC 59267-1404-02
• Does not include an ancillary kit

Pediatric Pfizer-BioNTech (Vaccine)
100-dose minimum order
• 10-dose multi-dose vial
• NDC 59267-1055-04

Pediatric Pfizer-BioNTech Bivalent Booster
100-dose minimum order
• 10-dose multi-dose vial
• NDC 59267-0565-02

Pediatric Pfizer-BioNTech (Vaccine)
100-dose minimum order
• 10-dose multi-dose vial
• NDC 59267-0078-04

Pediatric Pfizer-BioNTech Bivalent Booster
100-dose minimum order
• 10-dose multi-dose vial
• NDC 59267-0609-02

ARIZONA DEPARTMENT OF HEALTH SERVICES

COVID-19 Vaccine Presentations Available to Order in ASIIS

Moderna

Pediatric Moderna (Ages 6 months - 5 years) - Blue Cap with Magenta Border on label
100-dose minimum order
• 10-dose multi-dose vial
• NDC 80777-0279-99

Moderna Bivalent Booster (Ages 6+) - Blue Cap
100-dose minimum order
• 5-dose multi-dose vial
• NDC 80777-0282-99

Moderna Bivalent Booster (Ages 6 months - 5 years) - Dark Pink Cap with Yellow Border on label
20-dose minimum order
• 2-dose multi-dose vial
• NDC 80777-0283-99

Novavax

Novavax COVID-19 Vaccine, Adjuvanted
(Indicated for intramuscular injection)
100-dose minimum order
• 10-dose multi-dose vial
• NDC 80651-0102-10

Novavax (Ages 12+)
50-dose minimum order
• 5-dose multi-dose vial
• NDC 80651-0102-10

602-364-3899
ASIISHelpDesk@azdhs.gov
<https://azdhs.gov/vfc>

Updated 03/27/23

Pfizer pediatric doses (5-11) should NOT be reported as adult doses

Reminder:

The CPT and CVX codes for the Pfizer age 5-11 vaccine and the Pfizer age 12+ vaccine are different.

Please ensure that your EHR has been updated to document the vaccines correctly.

For information on Arizona HL7 Specific Rules for Version 2.5.1, click [here](#).

For information on List of Vaccine Names, Best ASIIS Selection and CPT/CVX Codes, click [here](#).

List of Vaccine Names, Best ASIIS Selection and CPT/CVX Codes



This list matches the vaccine name or codes in Arizona State Immunization Information System (ASIIS) with the brand name or other common names of the vaccines you use most often.

Vaccine trade name or common name	Fund	Best ASIIS Selection	Age (Range)	Dose	Route	Manufacturer/ NDC Number	CPT Code	CVX Code
COVID-19 Vaccines								
COMIRNATY Pfizer COVID-19 Vaccine	PAN	COVID-19, mRNA, LNP-S, PF, 30 mcg/0.3 mL dose	16+ years 12+ years EUA	0.3 mL	IM	Pfizer, INC - PFR NDC: 59267-1000-02 (195pack-dose vials) (1170 doses)	91300	208
	PAN	COVID-19, mRNA, LNP-S, PF, 10 mcg/0.2 mL dose, tris-sucrose	5-12 years EUA	0.2 mL	IM	Pfizer, INC - PFR NDC: 59267-1055-04 (10pack-10dose vials) (100 doses)	91307	218
Moderna COVID-19 Vaccine	PAN	COVID-19, mRNA, LNP-S, PF, ...	18+ years ...	0.5 mL	IM	Moderna - MOD NDC: 80777-0273-99 (10pack-10dose vials) (100 doses)	91301	207
Janssen COVID							91303	212
AstraZeneca COVID							91302	210
Novavax COVID-19								211
Sinopharm (BIBP) C								510
Coronavac (Sinovac)								511
SARS-CoV-2 (COVID								213

Arizona HL7 Specific Rules for Version 2.5.1

The following specifications and rules supersede CDC and general HL7 guidelines when sending messages to the Arizona Immunization Program. Otherwise CDC and general HL7 guidelines apply.

- MSH.1 – Field Separator will be the pipe | identified as (ASCII 24)
- MSH.2 – Encoding Characters will be ^~\& identified as (ASCII 94, ASCII 126, ASCII 92, ASCII 38) respectively.
- MSH.3 – Sending Application is **required**
- MSH.4 – Sending Facility is **required** and shall be the IRMS ID assigned and provided by the ASIIS System.
- MSH.5 – Receiving Application is **required** and shall always be ASIIS.
- MSH.6 – Receiving Facility is **required** and shall always be ASIIS.
- MSH.7 – Date/Time of Message is **required** and shall be in the following format (YYYYMMDDHHMMSS) and have a degree of precision to the minute generated.
- MSH.9 – Message Type is **required** and shall be VXU^V04^VXU_V04 for Unsolicited Vaccination Messages.
- MSH.10 – Message Control ID is **required** and shall be unique for each message attempt from a sending facility with a maximum of 20 characters.
- MSH.11 – Processing ID is **required** and because the ASIIS system has separate points for

Vaccine Transfer Process

- Both the sending and receiving providers will email data logger reports to ArizonaVFC@azdhs.gov
- Transfers must be approved in ASIIS prior to moving the doses
 - [Enter the transfer in ASIIS](#) prior to moving the doses
 - Information you will need for ASIIS
 1. The organization and facility sending the doses
 2. The organization and facility receiving the doses
 3. The quantity
 4. The lot number
 - When the doses arrive, mark them [“received” in ASIIS](#). Do not administer doses before “receiving” them in the ASIIS inventory
- **Only onboarded, active COVID-19 vaccine providers may receive COVID-19 vaccines**
- Follow the [USP transfer guidelines](#) when packing the doses for transfer
 - Once frozen doses have been thawed they cannot go back into a freezer
- Wherever the vaccines are, data loggers must be with them to monitor the temperatures
- The ancillary kit must also be transferred with the doses

Vaccine Wastage

- **If vaccine expires before you can use it:**
 1. Follow the instructions on this [job aid](#) to process wastage in ASIIS. Please note that COVID vaccines will not need to be returned
 2. Dispose of the vaccine
- Expired/wasted COVID-19 doses should be disposed of in a sharps container or per the hazardous waste policy in your office
- Tear off the vial labels or mark the identifying information (lot number, NDC number, etc) with a black marker prior to disposing
- Expired/wasted COVID-19 vaccines **DO NOT** go back to McKesson or Pfizer at this time
- [How to Account for Wasted/Expired Doses Job Aid](#)

Correcting Negative Doses in ASIIS

Reconciling ASIIS inventory

- Giving more doses than what is in the ASIIS inventory will make the inventory
- negative, which will mean you cannot submit inventory reconciliation
- Only if extracting extra doses from a vial
 - Ex. Extracting 15 doses from Moderna 14-dose vial
- AIPO Train module: Correcting Negative COVID-19 Doses in ASIIS

How to Run Reminder/Recall Report

- ASIIS Reminder/Recall Reports AIPO Train course
- Identify patients due/past due for vaccines
- Schedule patients
- Create letters, postcards, and mailing labels

ASIIS Reminder Recall Reports



If your ASIIS Inventory is not accurate

- Doses that have been administered to patients should not be removed from the Reconciliation page (Inventory) in ASIIS
- Troubleshoot why the doses did not decrement
 - [Register for Dose Accountability Webinar](#) in AIPO Train
 - Past Dose Accountability Webinar videos in course module
- [Job aid](#) to walk you through process of searching/adding/editing patient records manually in ASIIS

Mark doses as *Received* in VOMS 2.0 when the doses arrive

- Required regardless what system you use to document vaccine administration (EHR, VMS, ASIIS)
- When the doses arrive, mark them as **Received** in VOMS 2.0
 - Log into ASIIS
 - Go to Inventory Management> VOMS 2.0> Orders & Transfers > Click Receive button under Action header for the Inbound Order/Transfer
 - Verify the expiration date, lot number, and quantity of doses
 - Click the **Receive** button to receive the doses into the ASIIS Inventory
- After the doses are **Received** in VOMS 2.0, you may administer them
- If you administer doses before they are **Received** into the ASIIS inventory, they will not decrement from the ASIIS inventory

Vaccine Order Details

VACCINE	LOT #	EXPIRATION DATE	RECEIPT DATE	FUNDING SOURCE	ORDERED QUANTITY	APPROVED QUANTITY	SHIPPED QUANTITY	RECEIPT QUANTITY	ACTION
Coronavirus (SARS-CoV-2)(COVID-19)									
Covid-19, Mrna, Lnp-S, Pf, 3 Mcg/0.2 Ml Dose, Tris-Sucrose Pfizer 6m-4y Pediatric Covid-19 Vaccine PFR 100 pack - VIALS NDC: 59267-0078-04	<input type="text"/>	<input type="text" value="Exp Date"/>		PAN	800	800		<input type="text"/>	<div>ADJUST</div> <div>ADD LOT</div>

CANCEL

RECEIVE

Funding Source - PAN

- The funding source should be PAN.
- If you have an EHR, your EHR vendor should send the code VXC50. On the user interface side, you will select PAN when administering.
- If you select VFC, State, Private, or something else as the funding source, the doses will not decrement from the ASIIS inventory.

Communication Resources


Granicus Communications (Email Updates)

The BIZS sends granicus updates (email updates) on a variety of topics including COVID-19 vaccines, ASIIS, routine vaccines, etc. This library can be accessed at any time with the above link to view previously sent updates.

Ad Council COVID-19 Vaccination Campaign: It's Up to You

getvaccineanswers.org


- Campaign site for consumers
- Answers common questions
- Provides vaccine information
- Link to Spanish site - www.DeTiDepende.org
- [Creative materials](#) that can be shared/used by providers



COVID-19 Vaccine Confidence Toolkit for Rural Jurisdictions

- Print ads
- Posters
- Brochure
- Social media posts
- Online resource guide
- PDFs can be customized using Adobe Acrobat Pro

Learn. Understand. Decide



LEARN.
UNDERSTAND.
DECIDE.

COVID-19 vaccine communication toolkit

Customize the templates within the COVID-19 vaccine toolkit to easily develop internal and external communication materials!

With the support of the [Delta Regional Authority \(DRA\)](#) and the Health Resources Services Administration's [Federal Office of Rural Health Policy \(FORHP\)](#), the [Delta Region Community Health Systems Development \(DRCHSD\)](#)

Vaccine Confidence
[Preventative Screenings Campaign](#)
[Swing Bed Campaign](#)
[Telehealth Campaign](#)

Events
[DRCHSD 2020 Virtual Summit](#)
[Program Introduction and Overview of Application Process](#)
[Frequently Asked Questions \(FAQs\)](#)

Upcoming Events

UPCOMING WEBINAR
[2021 DRCHSD Program Introduction and Overview of Application Process](#)
June 24, 2021
This 2021 DRCHSD Program Introduction and Application Process webinar will share the program purpose, goals, successes, eligibility criteria, and available resources.

UPCOMING WEBINAR
[Recruitment for Retention Hierarchy - Unleashing Your Most Powerful Recruitment Tool: Culture](#)
July 1, 2021
This session will focus on examining the 3RNET Recruiting for Retention Hierarchy, identifying gaps in the Hierarchy for your organization, and discussing how culture can become a powerful recruitment tool in rural and underserved areas.



UPCOMING WEBINAR
[Building the Foundation of Your Recruitment Efforts](#)
July 8, 2021
This session will focus on identifying the four phases of the recruitment process, distinguishing how the recruiting process in rural areas is different than urban, and

ADHS COVID-19 Communication Resources

- Posters in English and Spanish (sample [here](#))
- Addressing Misinformation ([here](#))
- Answering Patient Questions ([here](#))
- What to Expect After Getting COVID Vaccine ([here](#))
- What to Expect at Your Vaccine Appointment ([here](#))
- Vaccination Quick Answers ([here](#))
- Answering Common Patient Questions ([here](#))

COVID-19 Vaccine - Providers' Toolkit

- Order Form to request materials for your facility





One question you are likely to get is: "How do we know if COVID-19 vaccines are safe?"

- EXPLAIN:**
- FDA carefully reviews all safety data from clinical trials.
 - FDA authorizes emergency vaccine use only when the expected benefits outweigh potential risks.
 - ACIP reviews safety data before recommending any vaccine for use.
 - FDA and CDC will continue to monitor the safety of COVID-19 vaccines to make sure even very rare side effects are identified.


For example, you can say:

"COVID-19 vaccines were tested in large clinical trials to make sure they meet safety standards. Many people were recruited to participate in these trials to see how the vaccines offer protection to people of different ages, races, and ethnicities, as well as those with different medical conditions."



HHS Video “Tell Me More About Vaccines”

The video answers commonly asked questions about the COVID-19 vaccine. The video shares why vaccines are so important and provides expert commentary and graphic illustration to help viewers understand the science of vaccine development. Tune in to hear from various experts, including Dr. Anthony Fauci (NIH), Dr. Stephen Hahn (FDA), and Dr. Robert Kadlec (ASPR), on the steps researchers and scientists are taking to develop a safe and effective vaccine. You are encouraged to add the video link to your website or promote it on social media.



FDA Emergency Use Authorization for Vaccines Explained

- What is an Emergency Use Authorization?
- Are the COVID-19 vaccines rigorously tested?
- What safety and effectiveness data are required to be submitted to FDA for an EUA request for a vaccine intended to prevent COVID-19?

CDC COVID-19 Vaccination Communication Toolkits

- For Medical Centers, Clinics, Pharmacies, and Clinicians
- For Healthcare Professionals and Pharmacists
- For LTCF Administrators and Leadership
- For Employers of Essential Workers
- For Staff of Organizations Serving Communities
- [Pediatric Healthcare Professionals COVID-19 Vaccination Toolkit](#)

The toolkit contains a variety of resources that you can use virtually or in person (with proper COVID-19 safety precautions):

- Posters
- FAQ
- Key messages
- Slide decks

Pediatric Healthcare Professionals COVID-19 Vaccination Toolkit

Updated May 15, 2021 [Print](#)



As parents' most trusted source of information on vaccines, pediatric healthcare professionals play a critical role in helping parents/guardians understand the importance of COVID-19 vaccination and assuring them that [COVID-19 vaccines are safe and effective](#).

Your strong recommendation is critical for vaccine acceptance. Tell parents/guardians how important COVID-19 vaccines are to protecting their children's health.

Remind parents that after their family is fully vaccinated against COVID-19, they may start to do some things they had stopped doing because of the pandemic.

Even if you are not administering COVID-19 vaccines, you can help parents/guardians feel confident in choosing to get their children vaccinated against COVID-19 by addressing their questions and assuring them of the safety and effectiveness of COVID-19 vaccines.

The materials on this page will help you share clear and accurate information about COVID-19 vaccines when starting or continuing conversations with parents/guardians, as well as information for those who choose not to vaccinate their child.

Only healthcare professionals enrolled as vaccination providers directly through a health practice or organization can legally store, handle, and administer COVID-19 vaccine in the United States. Learn more about becoming a COVID-19 Vaccination Provider: [How to Enroll as a COVID-19 Vaccination Provider](#).

COVID-19 Vaccination of Minors

Patients, parents, and guardians may have questions about consent for vaccination for minors recommended to receive a COVID-19 vaccine. Learn more at: [Pfizer-BioNTech COVID-19 Vaccine](#).

Provider Resources for Patient Conversations About COVID-19 Vaccines

- [How to talk to your patients about COVID-19 vaccination](#)

CDC's Key Points to Communicate to Your Patients:

- COVID-19 vaccines are [safe and effective](#).
- Individuals 12 and up are all eligible to [get a COVID-19 vaccination](#).
- There are many locations to [find a COVID-19 vaccine](#).
- Having some [side effects](#) after vaccination is normal.
- You are not fully vaccinated until 2 weeks after the 2nd dose of a two-dose vaccine or 2 weeks after a one-dose vaccine.

Recipient Education



Talking to Recipients about COVID-19 Vaccines

Many people have questions about the new coronavirus disease 2019 (COVID-19) vaccines. As vaccine recipients' most-trusted source of information on vaccines, you play a critical role in helping them understand the importance of COVID-19 vaccination, as well as if and when it is likely to be recommended for them.

The materials below include proven communication strategies and tips for effectively setting expectations and addressing questions from COVID-19 vaccine recipients.



Making a Strong Recommendation for COVID-19 Vaccination



Answering Vaccine Recipients' Questions



Understanding and Explaining COVID-19 Vaccines

Background and safety information for healthcare professionals and other vaccine providers, as well as tips for explaining the vaccines to patients.

- [mRNA COVID-19 Vaccines](#)
- [Viral Vector COVID-19 Vaccines](#)

CDC COVID-19 Vaccine Web Pages for the general public



- [Benefits of Getting a COVID-19 Vaccine](#)
- [How COVID-19 Vaccines Work](#)
- [Frequently Asked Questions about COVID-19 Vaccination](#)
- [Understanding mRNA vaccines](#)
- [What to Expect at Your Appointment to Get Vaccinated for COVID-19](#)
 - Translated in many languages
- [When You've Been Fully Vaccinated](#)
- [Webpage COVID-19 Vaccines for Children and Teens](#)

COVID-19 Vaccine:

Helps protect you from getting COVID-19

Get a COVID-19 vaccine, wear a mask, stay at least 6 feet apart, avoid crowds, and wash your hands to protect against COVID-19.

[QUESTIONS & ANSWERS](#)[WHEN YOU ARE FULLY VACCINATED](#)[FOR HEALTHCARE WORKERS](#)



Getting Ready for Your COVID-19 Vaccine

[What vaccines are recommended for use?](#)[Do they work?](#)[Is it safe?](#)[Are there side effects?](#)[What if I am pregnant or breastfeeding?](#)

[WHAT YOU NEED TO KNOW >](#)

How Do I Get a Vaccine?

Search vaccine providers near you

[VaccineFinder](#)

- OR -

Check your state or territory's health department

[Select State / Territory](#)

FAQs

FAQs

Q: If I am a VFC provider and already have an ASIIS PIN, do I still need to onboard?

- A: Yes. If a facility location wants to administer COVID-19 vaccines, they will need to onboard and sign the CDC COVID-19 Agreement. The VFC Program and the COVID-19 vaccine program are separate programs with separate signed Agreements.

Q: How do I order COVID-19 vaccines in ASIIS?

- A: Onboarded and active COVID-19 providers can follow the instructions in this [job aid on how to place orders](#)

Q: Where can I find these slides?

- A: These slides can be found at azdhs.gov/covid19vaccine on the Provider Resources page. You can also find a link to the slides and the most recent recording in the Pandemic Provider Weekly Brown Bag course in AIPO Train.

THANK YOU!



ARIZONA DEPARTMENT
OF HEALTH SERVICES

Health and Wellness for all Arizonans